








Pharmaceutical Organic Chemistry-1

Chapter-5: Aldehydes & Ketones


Aldehydes & Ketones

Common Classes of Carbonyl Compounds			
General Formula	Class	General Formula	Class
	Aldehydes 		Ketones 
	Acid Chlorides 		Carboxylic acids 
	Amides 		Esters 

Aldehydes & Ketones




- Carbon is ***sp*² hybridized**.
- **C=O bond** is shorter, stronger, and more polar than C=C bond in alkenes.





			<i>length</i>	<i>energy</i>
	ketone C=O bond		1.23 Å	178 kcal/mol (745 kJ/mol)
	alkene C=C bond		1.34 Å	146 kcal/mol (611 kJ/mol)

Structure of Aldehydes and Ketones

- **Aldehydes and ketones** are characterized by the presence of the carbonyl group.



- **Aldehydes** have at least one hydrogen atom attached to the carbonyl carbon atom.
The remaining group may be another hydrogen atom or any aliphatic or aromatic organic group.
- The **-CH=O group** characteristic of aldehydes is often called a formyl group.
- In **ketones**, the carbonyl carbon atom is connected to two other carbon atoms.







Nomenclature of Aldehydes

IUPAC System

- Aliphatic aldehydes* are named by dropping the suffix -e from the name of the hydrocarbon that has the same carbon skeleton as the aldehyde and replacing it with the suffix -al.




Alkane - e + al = Alkanal



Nomenclature of Aldehydes




IUPAC System

- **Substituted aldehydes**, we number the chain starting with the aldehyde carbon.
 - **-CHO group** is assigned the number **1 position**.
 - Aldehyde group has priority over a double bond or hydroxyl group.




- **Cyclic aldehydes**, the suffix **-carbaldehyde** is used.

Common
(IUPAC)



IUPAC System


- Aromatic aldehydes** are usually designated as derivatives of the simplest aromatic aldehyde, **benzaldehyde**.*




Nomenclature of Ketone

Common


- Common names of ketones are formed by adding the word *ketone* to the names of the alkyl or aryl groups attached to the carbonyl carbon. **Alkyl ketone.**
- In still other cases, traditional names are used.




Acetone
(Dimethyl ketone)




Ethyl methyl ketone




Methyl vinyl ketone



Cyclopentyl ethyl ketone



Methyl phenyl ketone
(Acetophenone)











Diphenyl ketone
(Benzophenone)

IUPAC

- In the IUPAC system, the ending for ketones is -one.
- The chain is numbered so that the carbonyl carbon has the lowest possible number.
- For cyclic ketones, numbering always starts from the C=O group.
- The prefix "oxo" is used when the ketone is not the principal functional group.

IUPAC
(Common)





Nomenclature of Aldehydes Ketones

NOTE

S In common names carbon atoms near the carbonyl group are often designated by Greek letters.


- The atom adjacent to the function is *alpha* (α), the next removed is *beta* (β) and so on. Since ketones have two sets of neighboring atoms one set is labeled α β etc., and the other α' β' etc.




- The functional group priority order in nomenclature system is as following:
Acid and derivatives > aldehyde > ketone > alcohol > amine > alkene > alkyne > ether

The Carbonyl Group

- **The structure and properties of the carbonyl group.**
 - The carbon–oxygen double bond consists of a sigma bond and a pi bond.
 - The carbon atom is sp^2 -hybridized. The three atoms attached to the carbonyl carbon lie in a plane with bond angles of 120° .
 - The pi bond is formed by overlap of a p orbital on carbon with an oxygen p orbital.
 - There are also two unshared electron pairs on the oxygen atom.
 - The C=O bond distance is $1.24\text{\AA}$ shorter than the C-O distance in alcohols and ethers ($1.43\text{\AA}$)




(a)




(b)

The Carbonyl Group

- Oxygen is much more electronegative than carbon. Therefore, the electrons in the C=O bond are attracted to the oxygen, producing a highly **polarized bond**.






- As a consequence of this polarization, *most carbonyl reactions involve nucleophilic attack at the carbonyl carbon*, often accompanied by addition of a proton to the oxygen (electron rich).




Physical Properties of Aldehydes and Ketones

Boiling Points

- Carbonyl compounds boil at higher temperatures than hydrocarbons, but at lower temperatures than alcohols of comparable molecular weight.



- This is due to the intermolecular forces of attraction, called dipole-dipole interactions, which is stronger than van der Waals attractions but not as strong as hydrogen bonds.




Dipole-dipole attractions among carbonyl compounds

Physical Properties of Aldehydes and Ketones

Solubility



y

- Carbonyl compounds as aldehydes and ketones have a **C=O bond**, but no **O-H bond**, cannot form hydrogen bonds with themselves.
- The **polarity of the carbonyl group** also affects the solubility properties of aldehydes and ketones.
- Carbonyl compounds with **low molecular weights** are soluble in water as **they can form hydrogen bonds** with O-H or N-H compounds.




1) Oxidation of Primary and Secondary Alcohols

- Oxidation of **secondary alcohols** yields **ketones**.





2) Hydration of Alkynes

- Hydration of **acetylene** yields acetaldehyde (catalyzed by acid and mercuric).






2) Hydration of Alkynes

- Hydration of **terminal alkynes EXCEPT acetylene** yields ketones (catalyzed by acid and mercuric).





3) Ozonolysis of Alkenes

Product (aldehyde or ketone) depends on the **structure of alkene**.



4) Friedel-Crafts Acylation


Preparing ketones that contain an aromatic ring.




Reactions of Aldehydes and Ketones

A) Reduction of Carbonyl Compounds

- Aldehydes and ketones are **easily reduced to primary and secondary alcohols**, respectively.
- The most common metal hydrides used to reduce carbonyl compounds are **lithium aluminum hydride (LiAlH_4)** and **sodium borohydride (NaBH_4)**.




- **Example:**





B) Oxidation of Carbonyl Compounds

- Oxidation of aldehydes gives a carboxylic acid with the same number of carbon atoms.
- Because the reaction occurs easily, many oxidizing agents, such as KMnO_4 , CrO_3 , Ag_2O and peracids (such as, perchloric acid HClO_4 , and permanganic acid HMnO_4). will work




- **Example:**



Reactions of Aldehydes and Ketones

C) Nucleophilic Addition Reactions

- Nucleophiles attack the carbon atom of a carbon-oxygen double bond because that **carbon has a partial positive charge**.
- The overall reaction involves addition of a **nucleophile and a proton across the pi** bond of the carbonyl group (when carried out in alcohol or water).





C) Nucleophilic Addition

Reactions

1) Addition of Grignard Reagents: Formation of

Alcohols Grignard reagents act as carbon nucleophiles toward carbonyl compounds.

- The reaction of a Grignard reagent with a carbonyl compound provides a useful route to alcohols.




- The type of carbonyl compound chosen determines the class of alcohol produced

C) Nucleophilic Addition


Reactions

1) Addition of Grignard Reagents: Formation of Alcohols


Formaldehyde gives primary alcohols.



○ *Other aldehydes give secondary alcohols*



○ *Ketones give tertiary alcohols.*




C) Nucleophilic Addition



Reactions

2) Addition of Hydrogen Cyanide: Formation of

Cyanohydrins
Hydrogen cyanide adds to the carbonyl group of aldehydes and ketones to form **cyanohydrins**, compounds with a hydroxyl and a cyano group attached to the same carbon.



○ Example



C) Nucleophilic Addition

Reactions

3) Addition of Ammonia and Ammonia

Derivatives The addition of nitrogen nucleophile, such as ammonia(NH₃) and substituted ammonia (NH₂-Y).

