The Lymphatic System

Dr. Mustafa Saad
(2021)

The Lymphatic System

Emain Component UI cacdiovax VII (i.e.) Lymphatic UI

System

Blood UI

Lymph JI

Carried lymph nodes

Lymphatics of the
marriary pland

Lymphatics of the
upper sinb

Hiss Me

Lissue

Lymphatics of the
upper sinb


UI sils UID

Lymph, Blood)

Thoracic duet

T

Lymphatic System Structure and Function



2

الكاستابة العناعية.

Lymph

- Most components of blood plasma filter through the capillary walls to form interstitial fluid.
 - More fluid filters out of blood capillaries than returns to them by reabsorption.

السائل التي تعن تعنية السائل التي الله التي تعنية تعنية الله The excess filtered fluid drains into lymphatic vessels and becomes lymph.




Fig.2: Capillaries and lymphatics.

وادال المil Fluid الى مهارالة Filtration والخنعلم reabsorption ووالخنعلم Blood الى Filtration وحفل دافل المناه المناه Lymph المناه Lymph المناه المن

Lymphatic Vessels

- ☐ Vessels begin as lymphatic capillaries These are closed at one end.
- Lymphatic capillaries unite to form large lymphatic vessels.

 These resemble veins in (محالات عليه المحالة structure but thinner walls and ومحالة ومحالة عليه المحالة المحالة
 - A series of swellings are present along the lymphatic vessels.

 These are the lymph nodes.




Fig.3: The distribution of lymphatic vessels in the body showing the lymph nodes. Also shown is the closed end of a lymphatic capillary.

عقدليعفاوية


Ultimately, the lymph drains into 2 main channels:

5-18/1/E)

Thoracic duct: a long duct that drains lymph from the entire left half of the body and the right half below the ribs including the right lower limb.

Right lymphatic duct: a short duct that drains lymph from the right side of the body above the ribs including the right upper limb.

These ducts open into the junction of the subclavian and internal jugular veins as they form the brachiocephalic veins; thus, lymph will return to the blood.






Fig.4: Above, areas of lymph drainage to the two main lymphatic ducts. Left, The opening of the ducts into the veins.

5

Lymphatic Organs and Tissues

- 2 groups based on function
 - Primary lymphatic organs:
 - Sites where B and T lymphocytes are formed and mature.

المحتمدة المحتمدة

Red bone marrow and thymus.

Secondary lymphatic organs

- Sites where most immune response occurs, meaning that they are the sites where the mature lymphocytes perform their function.
- Lymph nodes, spleen, lymphatic nodules.

الطحال

The Thymus→


lomphatic organs.

- The thymus is an asymmetric bilobed organ where mature Tat low pi cells are formed.
- النفف العلوي It's located in the superior mediastinum just behind the manubrium. It may descend into the anterior mediastinum to lie between the sternum and the pericardial sac. Sometimes it may ascend into the neck reaching as high as the thyroid gland.

JUSTE CAPSAIRS able Thymns JI Leave Capsute The fibrous capsule that surrounds the gland sends connective tissue trabeculae into the gland dividing it into lobules Each lobule is formed of a dark outer region with immature T-cells (the cortex) and an inner lighter region with mature T-cells (the

عنا في منطقة ال عادة عنا تفرعان هاي التفرعان عبارة عن منطقة ال معتدون العادم ومتون ومتون

Thymus le le le le gland supre et structure une de inner de 7



Left lung

Diaphragm

The thymic tissue is most abundant in younger ages As the person grows, this tissue is gradually replaced by fatty tissue.

Red Bone Marrow - Blood. Blood.

- Red bone marrow is the site of formation of the blood elements: red blood cells, white blood cells and platelets. مِمَا يُح وموية
- Inside the red bone marrow, B lymphocyte form and mature.

 T lymphocytes are formed in the red bone marrow but they're

 immature. T-cells become mature in the thymus.
 - Red bone marrow in adults is present in the flat bones and the epiphyses of some long bones.

9

secandary Lymph 11 Uplie

Lymph Nodes

حيكل صدة الفول/الفاصوليا و

- Lymph nodes are capsulated bean shaped structures that are found along the course of the lymphatic vessels. They're usually present in groups and they're scattered all over the body.
- The capsule sends trabeculae into the node dividing it into compartments. Beneath the capsule there's a space called the subcapsular sinus.
- The node has an outer cortex with lymphatic nodules formed of B-cells and plasma cells. Deep to it is the inner cortex formed of T-cells with no nodules. Deep to the cortex is the medulla which contains B-cells, plasma cells, and macrophages embedded in reticular fibers.
 - * Between the lymphatic tissue in the cortex and medulla are spaces called cortical and medullary sinuses.

- In any organ in the body, the tissues forming the organ can be divided into two types:
- 1. The Parenchyma: any tissue in the organ responsible for performing the function of the organ. Example: in the lymph nodes, the lymphoid tissue is the parenchyma.
- The Stroma: tissues in the organ that are not related to the function of the organ but play a more related. function of the organ but play a supporting role. Example: in the lymph nodes, the capsule, the trabeculae, the reticular fibers, and fibroblasts are the stroma.


11


محدب معلم (فارع ليرا)

61,00 From the convex side of the lymph node several incoming lymphatic vessels enter the node. These are called afferent lymphatic vessels.

عيير (دافل لحوا) The concave side of the node is called the hilum. From it, one or two outgoing lymphatic vessels leave the node. These are called efferent lymphatic vessels. Also through the hilum arteries and nerves enter and veins exit the node.

تدفق The flow of lymph in a lymph node





Functions of lymph nodes: → Filter of lymph

Antigens in the lymph are trapped and the lymphocytes in the nodes react to it and initiate the immune response.

Macrophages in the node Macrophages in the node may directly destroy the antigen. صو لد من

It's important to know what lymphatic vessels drain a certain organ and what are the lymph nodes in its course, because this represent a pathway by which infections and cancer cells can spread. ए । हा । हा ।

14

الحبر كتله من الأنسعة اللينادية في م The Spleen

Tylien Largest single mass of lymphatic tissue in the body. It's an oval, soft organ located in the left hypochondriac region 1/13/0/12/

■ Its superior surface is smooth and related to the diaphragm and ribs 2010 and 11. Its visceral surface is irregular and related to various organs.

Histologically formed of:

1. White pulp - lymphatic tissue (lymphocytes and macrophages) surrounds branches of splenic artery (the artery that supplies the المحمال الطعاليه spleen and enters through the hilum).

Red pulp - blood-filled venous sinuses surrounded by splenic cords which contain red blood cells, macrophages, lymphocytes, and plasma cells@

■ Functions: (1) Destruction of old red blood cells by macrophages, (2)Storage of blood, (3)Filter of blood, (4)Formation of blood cells during fetal life

الناء الحياة الحنسية

15




Fig.7: The spleen and its histology.

Lymphatic Nodules . Wo opinio lail Lymphatic us is a soles nodes nodes lymphatic us is a Lymphatic us capsule Lymphatic us is a Lymphatic usion nodes nodes

□ Collection of lymphatic tissue not surrounded by a capsule.

Scattered throughout lining of gastrointestinal, urinary, reproductive, and respiratory tracts.

الم Most of them are small and solitary.

□ Some are large – tonsils, Peyer's patches, appendix.

عبون عنا که لوزان که لوزان که لوزان که نوزان که کیم که کیم که کیم که کیم کارست کیم کارست کار