Skin and soft-tissue infections

Dr. Isra Dmour

Credit: Dr. Rawan Abudalo

1

* Skin infections may be cause by Virus, bacteria and fungal

Skin defense mechanisms A huge population of normal microflora colonizes on skin nutrients. Oil is decomposed by some M.O giving fatty acids that reduce the pH (5.5) which acts as a first line defense (innate defense) as many M.O can't grow at this pH Sweat is rich in salt, which acts as another defense. However, few M.O are salt tolerant تقدد خلايا المرة سية وهاد (4) Rapid shedding of skin cells. desse barrier dem Protective mechanisms Present Normal Microflora Sweat which Contain Salf that Is frevent Mo growth parity Mechanism sproperties acidic e Salt بعلم وليع ملية القالقيان بوسط مليء براهانقيان براهان براهانقيان براهانق Skin intections as

Normal flora of skin

Skin رخا infections تکون سبب Stathylaccol ولفلهالغالح

antistaph ?

The normal skin is colonised mainly by <u>Gram positive</u> microorganisms (staphylococci and diphtheroids).

- Some Gram-negatives and yeasts, which can survive in the relatively harsh conditions – dryness, acidity and the presence of fatty acids, sebum and salt.
- The normal skin flora protects against infection by other microorganisms that must compete to survive .

antcgorism قعللا مراجع


2

الطبعة الأولى من منع

Pathogenesis of skin

- Superficial skin infections affect the epidermis and dermis, e.g. impetigo
- Deeper infections affect the subcutaneous tissue, e.g. cellulitis
- Infections of skin structures cause folliculitis.

 In hair felicule
- Infections may arise from pathogens that gain entry through damaged skin (streptococcal cellulitis).
- Bacterial toxins produced by infections at other sites may also result in skin changes (group A streptococci causing scarlet fever)



Systemic justs

Systemic justs

Infection

Suso Skin cui

DISEASES OF THE SKIN

easy to be control and breated marks

threatening infection and many can be treated with empiric oral antibiotics, with occasional surgical drainage (folliculitis, acne, impetigo, and cellulitis). Severe infections require IV antibiotics.


2. Complicated disease: have a high risk for life- or limb-threatening infection, and often necessitate hospital admission, surgical debridement, along with broad-spectrum antibiotic therapy

المجابع تعمل مواحق مواحق

Bacterial Skin Diseases						
Type of infection	Appearance of Lesion	Description of Lesian	Layer of Skin Involved	Common Pathogens	Treatment	
Inceipo Contagious Life and		Veddles with honey colored crust, often on the face of a child	Epidernis Zeo	Staphylococcus purees, Staphylococcus pyrogenes	Few lesions, topical antibiotics (e.g., mupinocn); humerous lesions; systemic therapy (e.g., cephalicia; dindamycin)	
Celulos		Enthematus diffue. But lesson with tregular border Spacial definits Enthematus definits	م م م م م م م اد علية م	Street eventus progenes Simply or an application > Supply over an arrange	ed colymposition of control	
(Maults		localized inflament populars anisating a small amount of par inflamention of hair folicide	Har folicle	S arms Personals distributed assisted with his tuby	Antibulica other natures warm moles compresse are uteful	

Bacterial Skin Diseases

- Acne occurs in both males and females
- M.O feed on sebum.
- Ducts of glands & and tissue become inflamed
- lesions occur within sebaceous follicles with keratin plugs blocking the sebaceous canal, resulting in 'blackheads.
- Blackheads" are mild form of acne where hair follicles & sebaceous glands are plugged with sebum & keratin, as the surface oxidizes, they take black color.



Canal Espice Sebum ije gland lie to * inflammation duct is Sebum le معنی الفت کو استان المال و معلی و معلی

Bacterial Skin Diseases- Acne

- Bacteria, especially Propionibacterium acnes, infect the area and cause more inflammation, more tissue destruction, and scarring.
- · Acne is treated with:
- 1. frequent cleansing of the skin and ما العنول العنوال العنو
- 2. topical ointments to reduce the risk of infection
- · (Accutane®) Isotretinoin (Vit A derivative) for severe & persistent cases, it inhibits sebum production but has serious side effects

عداج فعوصات قبل أمنه ه مدينه ليشهات فلعية في الدّمنة ماريده لعده





ومعات

Bacterial Skin Diseases -Burn Infections

Burn Infections Severe burns destroy much of the body's protective
 Covering and provide ideal conditions for infection

• Burn infections (mainly nosocomial), are responsible for 80% of deaths among burn patients.

 Pseudomonas aeruginosa is the prime cause of life-threatening burn infections. Many strains of these Gram-negative bacilli are antibioticresistant.

Pseudomonas aeruginosa infection should be suspected when a greenish discoloration appears at the burn site and cultures have a grapelike odor.



0

dead cell (eschan) (crust) —

Upo exc homer USin six lin relation Pix +

Bacterial Skin Diseases -Burn Infections

 The thick crust formed over severe burn (eschar) lacks blood vessels & bacteria grow beneath it cause severe infections → antibiotics can not reach these areas,

Removal of eschar by surgical scraping (debridement) helps antibiotics reach these sites.



Bacterial Skin Diseases

- Associated with deep wounds, gas gangrene often is a mixed infection caused by two or more species of Clostridium, especially C. perfringens.
- In body regions where oxygen concentration is low, spores germinate, multiply, and produce toxins and enzymes such that kill other host cells and extend the anaerobic environment. المحمدة على المحمدة الم

enzyme toxin join a ciac considera enzyme toxin join a concept describit of the concept described and concept described and

11


Bacterial Skin Diseases

- Gas gangrene
- · Foul odor is such a prominent feature of gas gangrene
- Characterized by high fever, shock, tissue destruction & blackening of skin, if not treated causes death
- Treatment: Penicillin& removal of dead tissue or amputation to affected limbs
- Controversial treatment: use of hyperbaric O₂ (inhibits obligate anaerobes)
- Prevention: adequate cleaning of wounds, delaying closing of wounds & provide drainage when they are closed.

عرونفا لعواء (02) حتى يلتم الجرع المحالفة (02) حتى يلتم الجرع

12

العفر)



Viral Skin Diseases

Complecation

Measles, or German Measeales (rubeola)

If the virus invades the lungs, kidneys, or brain, the common childhood disease often is fatal.

- Transmission. The measles virus is highly contagious; its portal of entry is the respiratory tract while coughing and sneezing.
- Treatment: alleviating symptoms

Prevention: vaccination (MMR). Immunity acquired after the

disease is lifelong.

Cesp tract ع عطاس سعال عن طريق

One hoaix spi ése le vique apple apel è vien de le Long life immunity

(Symflomatic) 96

Control

Teratogenesis

Rubella

Rubella is caused by the Rubella virus that

invades the lymph nodes, eyes, and skin.

Usually mild, but can get severe in Spregnant women کان مناف

Measles is caused by Morbillivirus that infects the respiratory system.

Characterized by a pink rash that starts in the face and spreads to the body, lasting for three to five days. The rash may be itchy and associated with other symptoms, such as red and watery eyes, running nose, headache, and joint pain.

Severe and can be life-threatening.

Symptoms start with high fever, runny nose, cough, and red, watery eyes. Tiny white spots with bluish-white centers on a red background are found inside the mouth on the inner lining of the cheek (Koplik's spots) two-three days after the initial symptoms. The rash appears as flat red spots on the face at the hairline and spreads downward to the neck, trunk, arms, legs, and feet

اذا شخس عطاب وقع الخرج ما يستلط بالناس وسيخنا أوراب فاحه فيه وماسشاركها لانه هالمرض معدى جدا

Viral Skin Diseases

Complexations


Measles, or German Measeales (rubeola)

If the virus invades the lungs, kidneys, or brain, the common childhood disease often is fatal.

- · Transmission. The measles virus is highly contagious; its portal of entry is the respiratory tract while coughing and sneezing.
- Treatment: alleviating symptoms

Prevention: vaccination (MMR). Immunity acquired after the

disease is lifelong.



où e long life immunity (MMR) Long life immunity

Symflomatic) Control

Rubella

Rubella is caused by the Rubella virus that invades the lymph nodes, eyes, and skin.

Usually mild, but can get severe in pregnant women کانمن م يعل شوهات ملقبه TORSH Teratogenesis

Measles is caused by Morbillivirus that infects the respiratory system.

Severe and can be life-threatening.


Characterized by a pink rash that starts in the face and spreads to the body, lasting for three to five days. The rash may be itchy and associated with other symptoms, such as red and watery eyes. running nose, headache, and joint pain.

Symptoms start with high fever, runny nose, cough, and red, watery eyes. Tiny white spots with bluish-white centers on a red background are found inside the mouth on the inner lining of the cheek (Koplik's spots) two-three days after the initial symptoms. The rash appears as flat red spots on the face at the hairline and spreads downward to the neck, trunk, arms, legs, and feet

16

اذا شخص عماب فيهم كانم ما يختلط بالناس ويستخدا أورات خاصة فيه وعاليشاركها كانه هالمرض معدى عيدًا

ubject	Day	Date
4		
here Vin	15 C 1/25/19/11	رى بارغ لنوغا جشما +
•	70.00	وهر يوتر من
	The state of the s	वंद्रा त्यंश
Oci , Feve	- Stress qui Re	enfaction Jas
مناعة	Shig les	Infaction to zon i Ecos
gar, gar during an yang yan kang mengelang semang dan pendaman dalam dan		



Cervical Cancer

، يعالع ب OTC drugs الكاليال Viral Skin Diseases الكاليال الاظافر قاع القام وه-

Warts caused by human papilloma viruses (HPV) Human Papilloma virus-

HTV specifically attack skin and mucous membranes.

 Warts grow freely in many sites in the body -the skin, the genital and the oral cavity.

Sexual Trans 3 50

 The Nature of Warts. Warts vary in appearance, area of occurrence, and pathogenicity. Some are barely visible and self-limiting-that is, they do not grow or spread- and others, such as laryngeal warts, are larger but benign.

A few warts are malignant -cause 99% of all cases of cervical cancer



معكن يسفل فالمورف التي دام من الأروات مثل مقص الأطاف

(Freezing) ciresini ante lamo i Gan & Estell & (Illusts) Salicyclic - of

سبى كذا مع فاللس مول علمها عنى يعما من - Saligyclic acid

Viral Skin Diseases

Warts

Transmission. .. human papilloma viruses are transmitted by direct contact, usually between humans, or by fomites. - معال اودات فاصه

- · Dermal warts form when the virus enters the skin or mucous membranes through abrasions. Genital warts are sexually transmitted.
- Treatment:
- 1. cryotherapy (freezing & excision),
 - 2. caustic chemical agents,
 - surgery.





- The fungi that invade keratinized tissue are called dermatophytes
- fungal skin diseases are called dermatomycoses.

 They can be caused by any one of several organisms mainly: Epidermophyton, Microsporum, and Trichophyton.

· They cause various types of ringworm and attack the skin, nails, and hair. hain , Skin , nails , Scalp - target

Fungi that invade subcutaneous tissue live in soil or as airborne

They <u>enter</u> tissues through wounds

Subculoncoustori Fungal Jas tissue intections

A	Infection	Site	Dermatophyte	ngworm) infections Clinical features	
SCAP	Tinea corporis	Truck	Epidermophylon flocoosum	Pruritus; circular erythematous lesion, scaling, clearing from centre	
	trunk den	Limbs	Trichophyton spp. Microsporum spp.	and delivery	
	Tinea cruris	Groin	Epidermophyton flaccosum Trichophyton rubrum	Pruritus; lesions similar to tinea corporis	
القام الرياد athlets Fe	Tinea pedis	Feet	Epidermophyton floccosum Trichophyton spp.	Pruritus; cracking and scaling between toes	
	Tinea ungulum	Nais	Trichophytan spp.	Thick, yellowish nails with surrounding erythema	
up lea-	Tinea capitis	Hair	Trichophyton tonsurans	Erythernatous patches on scale with scaling: hairs break, leaving bald patches	
100	- he less	7	Microsporum audounii Microsporum canis	المعاى مش تعلية	_اقار جو
	alcodler Welsec		000	24	
C.	John Jak		مطلوب هاد الهدو	اد العامد ط مطلوب	

Common dermatophyte (ringworm) infections



Tinea capitis



Tinea pedis لم التهاب سن الميمايح

Tinea corporis

most Common Cause

Poor hygine backenal Jew infection infection as Secondary infection


Fungal skin diseases سِمِسِ نِمِهَ لَبِس


• Athlete's Foot. In athlete's foot, or tinea pedis عناماً عناماً عناماً المعالمات ال من الشراقارميكية معدد قيلي

- · Fungi invade the skin between the toes and cause dry, scaly lesions. Fluid filled lesions develop on moist, sweaty feet.
- · Then the skin cracks and peels, and a secondary bacterial infection leads to itchy, white areas between the toes.
- · Prevention of athlete's foot depends on maintaining healthy, clean, dry feet that can resist the opportunism of fungi.

Treatment: topical miconazole

Prevention: keeping clean & dry feet.





Fungal diseases Opportunistic fungal infections

Candidiasis

Transmited

disease

- Some fungi & yeast can invade human flature when immunity is impaired.
- Candida albicans (yeast) present as normal flora in GI و urogenital tracts. و عند المحادث ا
- Superficial candidiasis appears as thrush, (milky patches of inflammation on oral mucosa) in infants, diabetics, immunocompromised patients or those receiving AB for long time.

it causes <u>vaginitis</u> (in <u>pregnancy</u>, <u>diabetics</u>, those using contraceptives as sugar level is elevated in vagina).



oral candiasis

27

اکر النی الانتیا حساسه componises کے سنجاموں کے انکی الکی الکی قومنی کے Fungal infections ب

Fungal skin diseases


قشرة السعر

- Dandruff-flaking skin cells from an inflamed scalp.
- In dandruff sufferers, the cells flake off in big clusters in only 7 to 10 days
- The yeast, Malassezia, occurs in everyone's scalp, but only overgrows in some, causing inflammation, and excessive shedding.
- reatment: apply shampoos as ketoconazole.



23

* المواد الكمانية تزيد عددت القشرة



Infections of the eye

- Viral conjunctivitis:
- Viral conjunctivitis is commonly caused by adenovirus and is onen railed pink eye.
- It can also be caused by:

herpes simplex virus,

2. aricella zoster virus, and 3. various other viruses.



· Viral conjunctivitis often begins in one eye and then infects the other eye within watery eye Secreations, Italy, will be a few days.

Patients often present with an upper respiratory tract infection or recent ن لحدث سبت وجود exposure to someone with viral conjunctivitis. well the uffer Respiratory infection

 Antiviral medications may be an option if your doctor determines that your viral conjunctivitis is caused by the herpes simplex virus.

ointment

عمين الماد تا المقي بالملام الا

Contagiouse Infections of the eye المصاب العالة ماحدا ب

 Bacterial conjunctivitis, or pinkeye, is an inflammation of the conjunctive caused by organisms such as

- 1. Staphylococcus aureus,
- Streptococcus pneumoniae,
- 3. Pseudomonas species, and
- 4. Haemophilus influenzae.
- It is extremely contagious, especially among children.

- Topically applied sulfonamide ointment or erythromycin is an effective treatment.
- Do not share towels or eye cosmetics



Iliral Conjun is Su + (200) Watery Secretion ais 151 # كا اذا العن بنكن مسكرة ومَت الدستية الماستية المون النوع بكون Lo Bacterial Conju - Sticky Secretion can

- PATT TENEDERING