

Influenza Virus Infection عادة خلال الشتاء بسرعة يمكن بأي وقت

General Principles:

✓ Influenza is an acute febrile respiratory viral illness, readily transmissible and associated with outbreaks of varying severity during the winter months and with high mortality and high hospitalization rates.

✓ Seasonal influenza epidemics causes nearly 650,000 deaths each year globally, with the highest burden among children younger than 5 years and adults 75 years and older.


✓ Clinical presentation is similar to a number of other respiratory illnesses.

← نبأءا عليها

ببجل الينوصنا بسرعه

لانه سرعه نعيم بين و لين

other URI



ما يكون
قادر يعمل
His daily activities

عادة
يكون
بال upper
بسرعة
وعنه
ما ينزل
على ال lower
(Lungs)
يمكن
يعمل
primary
in influenza
pneumonia

✓ Severe illness in:

- older than age 65 years
- young children (< 2 years old) → حكمة من ال day cam
- underlying medical conditions (pregnancy and cardiopulmonary disorders)

immunocompromised, pregnant

✓ Incubation period 1 to 7 days. Adults are infectious from the day before their symptoms begin through 7 days after the onset of illness.

✓ Children can be infectious for longer than 10 days after the onset of illness.

✓ Influenza A and B viruses are the two types that cause disease in humans.

✓ Influenza A viruses are further categorized into different subtypes based on changes in two surface antigens—hemagglutinin and neuraminidase (NA).

✓ Influenza B viruses are not categorized into subtypes.

✓ Primary subtypes of influenza A (circulating among humans for the past 3 decades) are H3N2 & H1N1.

بعضهم
يعدوا من اليوم قبل ظهور
الأعراض لمدة تسع أيام
من ظهور الأعراض
(يعني من قبل ما تبدأ الأعراض)
يوم بدلتوا يعدوا

الأطفال
بعضهم
يعدوا فترة
أكثر من
الadult

(A)
كان يصيب
الحيوانات
وممكن

لنقل

من الحيوانات
للإنسان

inactivated
attenuated

Quadrivalent

حتى بناخذه (season)

صل هوو (3)

مطاعيم الانفلونزا

Combinational من ال Na/H
الي باخر 30 سنة
موجودين

بالثلاثي → فال Vassie
والثالث ← B
H3N2
H1N1

الرابع
H3N2
H1N2
2B

Influenza prevention

- Infection control measures (hand hygiene, basic respiratory etiquette (cover your cough and throw tissues away))
- Contact avoidance

Annual vaccination is recommended for:

- All persons age 6 months or older → من عمر 6 أشهر والأكبر
- Caregivers (eg, parents, teachers, babysitters) of children less than 6 months of age
- People who live with and/or care for people who are at high risk, including household contacts and healthcare workers.
- Pregnant women regardless of trimester (vaccination with IIV but not with LAIV). → inactivated vaccin
- Immunocompromised hosts should receive annual influenza vaccination (IIV but not LAIV)

preg
immuno
com → IIV (inactivated)
vaccines

IIV → inactivated influenza
vaccines
اختصار
virus

live attenuated
influenza vaccines

- Vaccine should be administered under the supervision of a health care provider who is able to recognize and manage severe allergic conditions (inpatient or outpatient medical setting).

[الاصلي عادي يعطي بس لازم رخصة]

- Ideal time: October/November (sufficient antibody titers after vaccination takes ~2 weeks).

لانه جهاز المناعة
حتى يعمل مناعة
والـ A B له
بده اسبوعين

- The specific strains included in the vaccine each year change based on antigenic drift.

point mutation

- LAIV should not be administered until 48 hours after influenza antiviral therapy has stopped and influenza antiviral drugs should not be administered for 2 weeks after the administration of LAIV because the antiviral drugs inhibit influenza virus replication.

LAIV ← في Live virus بس مضعف يعمل اعراض خفيفة

اذا امريض مائي Anti Viral لازم يحصل

العلاج وسني يومين عشان يوخد ال Vaccine

لو خلع ال Anti Viral ويغيب حوالي المطحوم رح يقضي على LAIV (لازم يستني يومين)

هون حتى لو اخذ المطحوم
وايضا Anti Viral غير بعد
2 weeks

Postexposure prophylaxis

- For seasonal prophylaxis and persons exposed to a household contact who were diagnosed with influenza.
- Antiviral drugs available for prophylaxis of influenza should be considered adjuncts but are not replacements for annual vaccination.
- Oseltamivir and zanamivir are effective prophylactic agents against influenza.
- Prophylaxis should be considered during influenza season for the following groups of patients:
 - Persons at high risk of serious illness and/or complications who cannot be vaccinated. →
 - Persons at high risk of serious illness and/or complications who are vaccinated after influenza activity has begun in their community.

مثلاً طفل
عاجل
7 أشهر

عد
immuno
corpsed
عنه
advanced
HIV

عد فتوقع
فايصير عنه
استفاده كانه
من الادوية

ارقطع
قبل بعد
10/11

oseltamivir → من
zanamivir → زعمه

Treat + prophylaxis → Antiviral
Mainstay prevention → Vaccines

Antiviral → بقطره
خلال 48
ساعة
من ابتداء مرض
الغايروسي

- Persons with severe immune deficiency or who may have an inadequate response to vaccination (e.g., advanced HIV disease, persons receiving immunosuppressive medications)- continued for

the duration that influenza viruses are circulating in the community during influenza season.

هناك الشخص لو اخذ الـ Vaccine خارج يكون ضاعه كافيه
بنحيه عن طريق انه لفظيه Antiviral طول فيه عايكون معرض

الموسم
الفاليردي

- After exposure to an infectious person (prophylaxis continued for 10 days after last exposure)

- Long-term care facility residents, regardless of vaccination status, when an outbreak has occurred in the institution.

Cancer Medication
Cyclosporin
Long-term Corticosteroid

حكت بي جيبها انه مريض

من حفس سنوآ عمل Kidney

Transplant

و فاني على Cyclosporin ولموره مقام

زي بدر رعايه كبار السن

لهون الجوان بي عطيه لـ Antiviral Medication

كسنان لم تعرضه لكون بيضاء خاصة لو بلس

لث في الالفريزا

هناك الرحي بعد

عاي تعرض للفايردي



لغي لو كنا

انه موسم او حذر بـ Antiviral
بالرعايه مريض بـ بـ بـ

Diagnosis:

✓ Clinical Presentation:

- Influenza virus causes an acute, self-limited febrile illness associated with rapid onset of fever, myalgia, headache, malaise, nonproductive cough, sore throat, and rhinitis.
- Nausea, vomiting, and otitis media are also commonly reported in children.
- Signs and symptoms typically resolve in 3 to 7 days. Cough & malaise may persist for > 2 weeks.



Otitis Media
Rhinosinusitis > complication
for influenza
خلال اسبوع ممكن
تضيق

عادة
حالتهم
الذين
يكون
Clinically
بس
هو
محب
هو
هل !!

✓ Diagnostic Testing:

Condition ~~respiratory~~ other over Lab مع مريضات over Lab في كين

- The sensitivity of clinical diagnosis ranges from 40% for children to 70% for adults and largely depends on the relative prevalence of influenza and other circulating respiratory viruses.
- Diagnosis is usually made clinically during influenza season, with confirmation by nasopharyngeal swab for rapid antigen testing, PCR (higher sensitivity), or direct fluorescent antibody test and culture.

Treatment:

- ✓ Treatment is usually symptomatic.
- ✓ Patients suffering from influenza should get adequate sleep and maintain a low level of activity.
- ✓ They should stay home from work and/or school in order to rest and prevent the spread of infection.

لـ زي تبع
Covid-19

→ شادراً
بذمعه

في ناس ممكن ابلش مع حتى لو بعد 48 ساعة
هذه الناس الي (critically ill) الي ممكن صابهم
و صاروا بار ICU

✓ Appropriate fluid intake should be maintained. Cough/throat lozenges, warm tea, or soup may help with symptom control (cough and sore throat).

✓ Antiviral medications may shorten the duration of illness but must be initiated within 24-48 hours of the onset of symptoms to be effective in immunocompetent patients.

✓ Antiviral therapy should not be withheld from patients presenting > 48 hours after symptom onset requiring hospitalization or at high risk for complications.

• The neuraminidase inhibitors (oseltamivir 75 mg PO q12h or zanamivir 10 mg inhaled q12h, each for 5 days, or peramivir, 600 mg single dose IV) are approved for the treatment of Influenza A and B.

• M2 inhibitors (amantadine and rimantadine, each 100 mg PO q12h) are not recommended owing to high rates of resistance.

فا بنبه ضم

X amantadine
rimantadine

ممكن
عند
asthma
HD
COPD
MI
IHD

دور
ال
انه يقل
لثة المرض

مجان

Oseltamivir
Zanamivir >

بقلوا
duration of
illness
by one day

of the onset of symp
بداء الاعراض
of patient presentation
لما تراجع الطبيب

[Anti Viral هو بديل غزار Vaccination]

- Circulating strains change annually with varying resistance patterns to both classes of antivirals. Treatment decisions must be based on annual resistance data, available from the Centers for Disease Control and Prevention (CDC) ([http:// www.cdc.gov](http://www.cdc.gov)).
- ✓ Vaccination is the most reliable prevention strategy.
- ✓ Annual vaccination is recommended for all individuals 6 months of age and older.
- ✓ Efficacy of vaccination varies annually from 50% to 90% depending on prevailing outbreak and circulating influenza strains.

← شركات الادوية لما بدوهم يعلو Vaccin هاي السنة
بشهر 15 او 9 يكون استقلوا عليه قبل بعدة اشهر
معناته ال strain اخدوها من تاعت الشتوية اطافية
فقدية الشبه بين ال strain تاعت الشتوية اطافية مع ال virus
تاعت صان السنة (بس رح يكون في Mismatch) عشتان

صية ال efficacy للطعيم من (50-90) % هو 100 %

الي علمهم لملاييت جدول التحديث
الجديدة الي نقتصرهم

ار كتاب انتشر ب 2020
والطاولات انظرهم 2020
في فصار
دراسات
رشدان
جديدة

diff
Mech
of
action
هذا صار
في دراسان
الخدم
2 antiviral
Medication
for
with
diff
MOA
for
influenza

TABLE 127-6 Recommended Daily Dosage of Influenza Antiviral Medications for Treatment and Prophylaxis—United States^{33,34,53}

Drug	Adult Treatment	Adult Prophylaxis*	Pediatric Treatment	Pediatric Prophylaxis*
CAP-dependent endonuclease Inhibitor				
Baloxavir ^{1,2}	12 yrs and older: 40 to <80 kg: One 40 mg dose >80 kg: One 80 mg dose	None حاليا صوة approved for prophylaxis	FDA approved and recommended for use in children 12 yrs or older weighing at least 40 kg. See adult dosage	None
Neuraminidase Inhibitors				
Oseltamivir ^{1,2}	75-mg capsule twice daily x 5 days	75-mg capsule daily x 10 days	Term Infants 0–8 months: 3 mg/kg/dose twice daily 9–11 months: 3.5 mg/kg/dose twice daily or 3 mg/kg/dose twice daily ≥1 year: ≤15 kg: 30 mg twice daily >15–23 kg: 45 mg twice daily >23–40 kg: 60 mg twice daily >40 kg: 75 mg twice daily Duration: All for 5 days	Not recommended if <3 months 3–<12 months, 3 mg/kg/dose daily 9–11 months, 3.5 mg/kg/dose daily ≥1 year: ≤15 kg: 30 mg daily >15–23 kg: 45 mg daily >23–40 kg: 60 mg daily >40 kg: 75 mg daily Duration: All for 10 days
Zanamivir	10 mg (2 of 5 mg Inhalations) twice daily x 5 days	10 mg (2 of 5 mg Inhalations) daily x 10 days	10 mg (2 of 5 mg Inhalations) twice daily x 5 days for ≥7 years old	10 mg (2 of 5 mg Inhalations) daily for ≥5 years old x 10 days
Peramivir ¹	13 yrs and older: One 600 mg dose via Intravenous Infusion for 15–30 minutes	None	2 to 12 yrs of age: One 12 mg/kg dose, up to 600 mg maximum, via Intravenous Infusion for a minimum of 15–30 minutes	None

antiviral → resistance
لما اكثر من ال AB

Beloxavir → Has Many ddt
هو لا complicated
influenza

صارت
من 6 اشهر
فو سنبين

يكن
ببفدر
نخطيه
من عمر
6 اشهر

Continued for Table 127-6

a If influenza vaccine is administered, prophylaxis can generally be stopped 14 days after vaccination for noninstitutionalized persons. When prophylaxis is being administered following an exposure, prophylaxis should be continued for 10 days after the last exposure. In persons at high risk for complications from influenza for whom vaccination is contraindicated or expected to be ineffective, chemoprophylaxis should be continued for the duration that influenza viruses are circulating in the community during influenza season.

b Time to peak = 4 hours. Food and cations (calcium, aluminum, magnesium, iron) can decrease peak concentration by 48%. Long half-life (79.1 hours) and is metabolized by UDP-glucuronosyltransferase (UGT1A3) and CYP3A4.

c For the treatment of uncomplicated influenza with oral baloxavir or intravenous peramivir, a single dose is recommended. Longer daily dosing (oral oseltamivir or intravenous peramivir) can be considered for patients who remain severely ill after 5 days of treatment.

d Oseltamivir dosing for preterm infants using their postmenstrual age (i.e., gestational age + chronological age): <38 weeks: 1.0 mg/kg/dose twice daily; 38–40 weeks: 1.5 mg/kg/dose twice daily; >40 weeks: 3.0 mg/kg/dose twice daily.

e In patients with renal insufficiency, the dose should be adjusted on the basis of creatinine clearance. See <https://www.cdc.gov/flu/professionals/antivirals/summaryclinicians.htm>.

f Some experts recommend 150 mg twice daily for severe illness in pregnant women. Optimal dosing for prophylaxis in pregnant women is unknown.

g The American Academy of Pediatrics recommends 3.5 mg/kg per dose twice daily; CDC and US Food and Drug Administration (FDA)-approved dosing is 3 mg/kg per dose twice daily for children aged 9–11 months.

Note: Although amantadine and rimantadine have been used historically for the treatment and prophylaxis of influenza A viruses, due to high resistance, the CDC no longer recommends the use of these agents for the treatment and/or prophylaxis of influenza

Beloxavir
سَيَّارٌ بِالْأَكْلِ
والإماتة

سَيَّارٌ ال
Peak
تبعه 48%

لو بده رجي
عليها Case

انه حد
باخذ حليب
يشرب كاسة

حليب او باصر

VitC - mg - potassium
او Calcium - iron - antacid
زيهيك ← كلهم باثرا عديم
لدرهم فانه يباع لينهم

مو موجودة
هناي السلايه ملكي صحتها
بالرئود على التي بس شرحها
بالمحاضرة يعني . سهله فابدها في

Complications:

- ✓ People at greater risk of complications are:
 - Adults > 65 years old
 - residents of nursing homes and other long-term care facilities
 - pregnant women (and those up to 2 weeks postpartum)
 - patients with chronic medical conditions (e.g., pulmonary disease, cardiovascular disease, active malignancy, diabetes mellitus, chronic renal insufficiency, chronic liver disease, immunosuppression including HIV and transplantation, morbid obesity)
- ✓ Influenza pneumonia and secondary bacterial pneumonia, typically due to *S. aureus*, are the most common complications of influenza infection.
- ✓ Viral antigenic drift and shift can cause emergence of strains with enhanced virulence or the potential for pandemic spread, requiring modified therapy or heightened infection control measures.

الهد فلكاوي
ادعولي لو استخذتوا عن الدفريغ
😊