

#### Physical Pharmacy I Lecture 1: Intermolecular forces

Dr. Isra Dmour

Credits: Dr. Dr. Areen Alshweiat, Nizar Al-Zoubi

Areen.alshweiat@hu.edu.jo


#### Intermolecular forces

- Intermolecular Vs. Intramolecular forces.
- Intermolecular forces: between molecules (e.g. Van der waal forces).
- Intramolecular forces: inside the molecule, i.e., between atoms or groups inside the same molecule (e.g., covalent, ionic, and metallic).

  The key difference between the two types is that the covalency is not established in intermolecular interactions  $\rightarrow$  less energy is required

to break intermolecular forces as compared to intramolecular ones.

(intra) II-



# Repulsive and Attractive Forces

Different charges

Similar charges = repulsion

عشاق بعد الله المحتمد على and the system is most stable.


والعزا الروا الروا


When two atoms or molecules are brought closer together, the opposite charges and binding forces in the two molecules are closer together than the similar charges and forces, causing the molecules to attract one another.


When the molecules are brought so close that the similar charges touch, they repel each other like rigid elastic bodies.)

At a certain equilibrium distance, (about 3-4 Å), the repulsive and attractive forces are equal. At this position, the potential energy of the


energy 11: 5 sels two molecules is a minimum







#### Repulsive and Attractive Forces



#### Intermolecular forces

Intermolecular forces can be divided into:

- · Van der Waals forces
- Ion—dipole interaction
- · Ion-induced dipole interaction
- Ion-ion interaction
- Hydrogen bonds

Ion: full charge eg: Nach (full charge) (Nat / Cb-1)
\*Water - hydrogen is partial Positive.

2 atoms lie is the Lils

is since of the more electronegative
than other

# Pole: 169 Polarity

incar had we had and and

. He had sendence he a day of the

المعاكث) (بسمب الع أكثر) المعاندة المعا

One of the atom in the covalent bond has the ability to attract the shared electrons to itself resulting in a polar molecules

• Difference in the electronegativity +

 dipole is a separation of two opposing charges over a distance r and generally described by a vector known as a dipole moment (μ).

التوزي والتحل المراج The symmetry of the molecules affects generally its dipole moment

(C) I in visus (O)

(C) I in visus (O)

(C) I in visus (O)

(Gas) a sublit HCL is I

ais all la

- · Van der Waal interactions are weak forces that involve the dispersion of charge across a molecule called a dipole.
- · Van der Waal interactions can be classified into:
  - A.Dipole-dipole interaction, orientation effect, or Keesom force

(تعفير عفين (تعفير عفين) عند (B)Dipole-induced dipole interaction, induction effect, or Debye force

C. Induced dipole-induced dipole interaction, dispersion effect, or London force

\* Dipole : net result and charges Positive, Negative.

#### Van der Waals forces

- The Partial opposite charges (permanent dipoles) attract one another (dipole-dipole interactions)
- Two polar molecules (two permanent dipoles)

The dipole-dipole forces increases as the polarity of the molecule increases.

\* electronegativity

(حدل الماية والموالية الموالية الموال

$$\delta + \delta - \delta + \delta - H - CI - H - CI$$

(Covelent not Ionic)

• Keesom forces occur between polar molecules in which the permanent dipoles interact with one another (dipole-dipole interactions) or (orientation effect).

Polar molecules have polar covalent bonds which are unevenly distributed in space due to the difference in the electronegativity of the atoms forming the bond e.g. HCl.


• The nucleus of the chlorine atom pulls the electron pair involved in the chlorine-hydrogen bond closer to itself and creates a permanent partial positive charge on the hydrogen and a permanent partial negative charge on the chlorine (Permanent dipole)

Dipole Moment has a
Magnitude and a Direction

## Van der Waals forces

Debye forces occur between a polar and a nonpolar molecule in which the permanent dipole in the polar molecule induce an electric dipole in the non polar one (dipole-induced dipole interactions) or (induction effect) e.g. water and oxygen.






• Example of Keesom forces

· (Acetone) Polar salvant

الأكسجين بتفل تسحب ع من الكربون (Dipol - Pipol)

## Van der Waals forces

- The oxygen molecule is nonpolar. (No charge)
- However, when it comes close to the oxygen atom in a water molecule, the partial negative charge on the oxygen atom repels the electrons in the oxygen molecule.
- This causes (induces) a temporary partial positive charge in the end closest to the water molecule and a buildup of a partial negative charge in the end furthest away. This induced dipole is temporary and forms only when the two molecules are extremely close to one another.




all istancell like (note)

Tils hold Induction I!

(Induction) me lo au lie

• This attraction is produced by temporary (asymmetry) in the distribution of the electrons around the nucleus.

الانكترونان سورة حول النواق على شكل (Orbitals) مزان



201.

#### Van der Waals forces

• The strength of Debye forces increases with the ease والم على كانت الروابط و of (distortion) of the electron cloud of the nonpolar معالم المستحون المتحريك المعادة المستحون المتحريك المعادة المستحون المتحريك المتحريك


• Debye forces is weaker than Keesom forces because the dipole in the nonpolar molecule is temporary (induced) and forms only when the two molecules are extremely close to each other.

Induction Amer relais

# Induction der Waals forces

London forces occur between two nonpolar weakest (neutral) molecules in which molecules can induce (neutral) molecules in which molecules can induce polarity on each other (induced dipole-induced dipole interactions) or (dispersion effect).e.g. (Helium)

Nobel gas Neon, hydrocarbons (Benzene, hexane, etc)...



#### Van der Waals forces

· London forces occur between all atoms and molecules (between polar/polar and polar/nonpolar molecules as well)

الفرو الدري بعس في ع محكم عليها فيحس جمها. The larger the atom or molecule, the greater its المدارات برداد (easier to induce a momentary dipole) and the عدد الدع فبرداد الما interaction forces become.

تنكوبا المسروحين. This is because:

- 1. The electrons are farther from the positive nucleus and so are held less strongly
- 2. The number of electrons is greater
- London Force is the weakest of all the intermolecular forces.

وكل ما كبر يهسي

جواً فريبة من النواه

Induction


- James


قوم معدى يوكي ل

Nacl - water

Ion-dipole forces


• These types of interactions account in part for the solubility of ionic crystalline substances in water; the cation, for example, attracts the relatively negative oxygen atom of water and the anion attracts the hydrogen atoms of the dipolar water molecules. > Negative






# Ion-dipole forces

• Ion-dipole forces occur between a charged ion and a polar molecule (i.e. a molecule with a dipole) Cations are attracted to the negative end of a dipole, while anions are attracted to the positive end of a dipole






20


# Ion-Induced dipole

- (Ion-Induced dipole forces occur between a charged ion and a nonpolar molecule.
- These forces result when the approach of an ion induces a dipole in an atom or in a nonpolar molecule by disturbing the arrangement of electrons






Spherical atom with no dipole. The dot indicates the location of the nucleus.



 $I_2+K+I^- = (K+I_3)$  net charge = Upon approach of a charged ion, electrons in the atom respond and the atom develops a dipole.


Non-Polar - Iodine(Iz) the atom develops a dipole.

(Ion) - Pottasium Iodide(KI)



#### Ion-ion interactions


- A cation on one compound will interact with an anion of another compound.
- Ion-ion interactions may be <u>intermolecular</u> (e.g. a hydrochloride salt of a drug) or <u>intramolecular</u> (e.g., a salt bridge interaction between counter ions in proteins).



المحقام أحوتنا عبارة عن Weak acids or bases فيعمال هانكه والماء بعنوله عليه عليه عليه عليه وعشان ندوبه بالماء بعنوله المحالي فيتزيد (Saft) لكاله بعيد دالهاء بالماء مالهاء مالهاء مالهاء مالهاء مالهاء مالهاء مالهاء والمعمودة

- Hydrogen bond

   Hydrogen bond is a strong type of dipole-dipole interaction that occurs between a molecule containing a hydrogen atom and a strongly electronegative atom such as fluorine, oxygen, or nitrogen
  - In order to create the bond, the hydrogen atom must be covalently attached to another electronegative atom.
  - A perfect example of hydrogen bond is water.
  - · Hydrogen bonds can also exist between alcohol molecules, carboxylic acids, aldehydes, and DNA and polypeptides > Proficers (a-a)



## Hydrogen bond

Hydrogen bond

الحاء الفرعادية Hydrogen bonds are responsible for many unusual physical properties of water including its abnormally low vapor pressure, high boiling point, and the greater

showledge le (volume of ice water.) (ice has large volume than liquid)

• Hydrogen bonding is stronger than all Van der Waals intermolecular forces (they are given their own classification), but are still weaker than (ionic) and covalent bonds. afull ionic

(hydrophobic) air lell eil eil ail (hydrophobic) air eil ell ail فجزيئات الماء بتذجذب نو المركز فالوارط بين الماء نفسه water+ air 11 in interaction 11 incs 95 i

## Bond energies

• Bond energy is a measure of the strength of bond.

Intramolecular forces: Ionic>covalent>metallic Intermolecular forces:

Reviser (Ton)

• Ion-dipole>ion-induced dipole>hydrogen bond>keesom>Debye>London

Importance of intermolecular forces

These forces govern the interactions between all kinds of matter (e.g., atoms, molecules, bulk)

They are important to understand the properties of gases, liquids and solids.

Understand various physicochemical phenomenon and processes:

repulsion agin is 50% ye

Interfacial phenomena: are the ones that occur at the limit between two immiscible phases, so-called surface or interface.

-(Flocculation of suspensions: is a process wherein colloids come out of suspension in the form of floc or flake. (not cake)

- Stabilizations of emulsions, Emulsion stability refers to the ability of an emulsion to resist change in its properties over time.

Anna Mail who will bit of

Remitader & emulsion: Avoid freezing

forces rolling is properties of le doselle sit alle per pair

Compaction of powders in capsules.

- Compaction of powders in capsules.

- Dispersions of powders or liquid droplets in aerosols

(An aerosol is a colloid of fine solid particles or liquid droplets, in air or another gas).

- Compression of granules to form tablets.

- Compression of granules to form tablets.

- Various intermolecular and intramolecular forces gover the secondary to the seconda

- Various intermolecular and intramolecular forces govern the secondary, tertiary and quaternary structures of biomolecules (e.g., proteins, DNA, RNA).

- They influence the stability, formulation and storage of pharmaceutical product derived from such biomolecules.

on details 11 عطوية حون