



لجان الترقيات



STATISTICS



MORPHINE ACADEMY

MORPHINE
ACADEMY

Hypothesis Testing



Hypothesis Testing

الهدف هو استبعاد احتمال أن تكون النتائج التي تم الحصول عليها من عينة ما ناجمة فقط عن الصدفة أو خطأ الحينة (sampling er.) وذلك بناءً على تفسير منطقي وهو ثوق لنتائج البحث

- The general goal of a hypothesis test is to rule out chance (sampling error) as a plausible explanation for the results from a research study.
- Hypothesis testing is a technique to help determining whether a hypothesis is true (e.g. treatment, procedure has an effect in a population), or simply if a relationship exists between two or more variables.

باختصار، اختبار الفرضيات يساعد في اتخاذ قرار مستند إلى البيانات حول صحة فرضية معينة.

Research Hypothesis vs Null Hypothesis

Research hypothesis (H_1) is what the research believes to be a true reflection on the general population. In another word, a true explanation for a phenomena in the population. The researcher wants to prove that his sample statistics is different than the population parameters. Research hypothesis is also called alternative hypothesis.
انه الباحث بدويثبت انه
إحصائيات العينة تختلف فعلياً
عن popu. parame.
مرات بجعلها (البديلة)

Null hypothesis (H_0) is the opposite of H_1 . The H_0 assumes no difference of test statistics and the population parameter. This means that the researcher hypothesis about a certain phenomena is not correct, and that there is no real difference between sample and population for a certain feature or difference is due another reason (that is not tested).
العكس تماماً
تفترض عدم وجود فرق، وأي اختلاف بسبب الاهدفة أو عوامل أخرى

Null & Alternative Hypothesis

Null Hypothesis or the Hypothesis to be Tested (H_0)

- A claim that there is NO difference between the population parameter like mean (μ) and the hypothesized value. In the testing process the null hypothesis either is rejected or not rejected (we do not say accepted).
→ مالتصلا
بشكل قاطع
رفض
عدم رفض

Alternative Hypothesis or the Research Hypothesis (H_A or H_1)

- A statement of what we will believe is true if our sample data cause us to reject the null hypothesis. Usually the alternative hypothesis is the research hypothesis (the conclusion that the researcher is seeking to reach).
← يوجد فرق حقيقي والباحث يسعى للثبات بناءً على البيانات

Null Hypothesis as an Assumption to be Challenged

- We might begin with a belief or assumption that a statement about the value of a population parameter is true.
- We then using a hypothesis test to challenge the assumption and determine if there is statistical evidence to conclude that the assumption is incorrect.
إذاً دليل قوي ضد الفرضية الصفرية لنستج انها غير صحيحة
- In these situations, it is helpful to develop the null hypothesis first.

باختصار:

- الفرضية الصفرية (H_0) هي نقطة انطلاق للاختبار.
- نختبر صحة H_0 بناءً على بيانات العينة.
- إذا وجدنا أدلة كافية، نرفض H_0 ، وإذا لم توجد أدلة كافية، لا نرفضها (لكن لا نقبلها بشكل قاطع).

q3.2

Example

- A new drug is developed with the goal of lowering blood pressure more than the existing drug.

تم تطوير دواء جديد

حفظ ضغط الدم

الحصول من الدواء
الموجود حالياً

- **Null hypothesis** :-

- The new drug does not lower BP more than the existing drug.

- **Alternative hypothesis** :-

- The new drug lowers blood pressure more than the existing drug.

شرح 3-

- الفرضية الصفرية تفترض عدم وجود فرق أو تأثير جديد (الدواء الجديد ليس أفضل من القديم).
- الفرضية البديلة تفترض وجود فرق أو تأثير (الدواء الجديد أفضل ويخفض ضغط الدم أكثر)

Null & Alternative Hypothesis

Gabapentin has no pharmacological effect

- The mean for population A is 20 ($H_0: \mu = 20$)
- The mean for population A is less than 20 ($H_0: \mu \leq 20$)
- The mean for population A is larger than 20 ($H_0: \mu \geq 20$)

Gabapentin has a pharmacological effect

- The mean for population A is not 20 ($H_1: \mu \neq 20$)
- The mean for population A is larger than 20 ($H_1: \mu > 20$)
- The mean for population A is less than 20 ($H_1: \mu < 20$)

هل جيب
Accepting or rejecting a hypothesis is not a proof of the hypothesis!
Null hypothesis can be true or false, we only can reject it or not to reject it

Null Hypothesis **vs** Alternative Hypothesis

Null hypothesis H_0	Alternative Hypothesis H_1
There is <u>no</u> relationship or difference	There <u>is a</u> relationship or difference
Refers to the <u>population</u>	Refers to the <u>examined</u> sample
Research aims to <u>reject</u> the <u>null</u>	Research aims to <u>accept</u> the <u>alternative</u>
Represent an original assumption	Prove statistically a systemic difference or relationship
Assumes a difference is due to chance	Assumes that difference is less likely to be due to chance.

Types of Statistical Errors in Hypotheses Testing

Because hypothesis tests are based on sample data, we must allow for the possibility errors.

✿ Type I error

- A type I error is rejecting H_0 when it is true.
- The probability of making a Type I error when the null hypothesis is true as an equality is called the level of significance (α) .

✿ Type II error

- A type II error is accepting H_0 when it is false.
- It is difficult to control for the probability of making a type II error (β) .
- Statisticians avoid the risk of making a type II error by using “do not reject H_0 ” and not “accept H_0 ”

Errors in hypothesis testing

TRUTH

DECISION

	Ho is true ($A=B$)	Ho is false ($A \neq B$)
Reject Ho ($A \neq B$)	Type I error “giving a treatment that does not work”	Correct 
Do not reject Ho ($A=B$)	Correct 	Type II error “not giving a treatment that works”

كيف أصبح فرضية جديدة

How to establish a good hypothesis?

تكون جملة واضحة

- Clear and declarative statement. Not a question.

ما تكون سؤال

- Show a relationship between variables

يجب ان يتم تحديد
كيف يؤثر متغير على
آخر أو العلاقة بينهما

- Reflect a body of literature or a theory

مباشرة و واضحة وغير مبهمة

- Be direct, explicit, and to the point.

لازم تكون الفرضية
مبنية على أساس علمي

- Be testable and measurable.

قابلية للاختبار
والقياس

Critical Value Approach to Hypotheses Testing

- **Step 1:** develop the null and alternative hypotheses.
- **Step 2:** specify the level of significance α . Common value of α are: $\alpha = 1\%$ or 0.01 , 5% or 0.05 , 10% or 0.1 هنا القسم الشائعة
- **Step 3:** collect the sample data and compute the value of the test statistics Z or t. جمع بيانات العينة
- **Step 4:** use the level of significance (α) to determine the critical value (tabulated value). لنستخدمها لتحديد القيمة
- **Step 5:** use the suitable rejection rule. الدرجة من الجدول الاحصائية
- **Step 6:** state the appropriate conclusions. ملخص

- تبدأ بتحديد الفرضيات ومستوى الدلالة.
- تجمع البيانات وتحسب إحصائية الاختبار.
- تحدد القيمة الحرجة بناءً على α .
- تقارن إحصائية الاختبار بالقيمة الحرجة.
- تقرر قبول أو رفض H_0 .
- تسجل الاستنتاج النهائي.

Step 1

- Set your hypothesis

Set both H_0 & H_1 .

Example

It is believed that a candy machine makes chocolate bars that are on average 5g. A worker claims that the machine after maintenance no longer makes 5g bars. Write H_0 and H_a .

Answer:

$$H_0: \mu = 5$$

$$H_1: \mu \neq 5$$

A company has stated that their straw machine makes straws that are 4 mm diameter. A worker believes the machine no longer makes straws of this size and samples 100 straws to perform a hypothesis test with 99% confidence.


Answer:

$$H_0: \mu = 4$$

$$H_1: \mu \neq 4$$


Step 2

- Set level of significance associated with the hypothesis.



Step 3


- Set the critical value needed to reject the null hypothesis (from Tables).
- You might need the table of Z-test, t-test, F-test.
- Based on the chosen table, look for the cut off value based on the level of significance you determined in step 2.



Right-tailed tests

$$H_0: \mu = 100$$

$$H_1: \mu > 100$$



Left-tailed tests


$$H_0: \mu = 100$$

$$H_1: \mu < 100$$

Points Left



alpha



100

Z_{crit}

Values that
differ “significantly”
from 100

Two-tailed hypothesis testing

- H_A is that μ is either greater or less than μ_{H_0}

$$H_A: \mu \neq \mu_{H_0}$$


- α is divided equally between the two tails of the critical region

- في اختبار ذو ذيلين، نبحث عن اختلاف في أي اتجاه (أعلى أو أقل) عن القيمة المفترضة μ_{H_0} .
- منطقة الرفض تقع في الطرفين (الذيلين) للتوزيع الاحتمالي.
- هذا يعني أنه إذا كان الإحصاء الاختباري يقع في أي من الذيلين (أقل من الحد الأدنى أو أكبر من الحد الأعلى للقيمة الحرجة)، نرفض الفرضية الصفرية.

Two-tailed hypothesis testing

$$H_0: \mu = 100$$

$$H_1: \mu \neq 100$$



One tale critical values

alpha .05, $Z_{\text{crit}}=1.64$;

alpha .01, $Z_{\text{crit}}=2.33$

Two tale critical values

alpha .05, $Z_{\text{crit}}=1.96$;

alpha .01, $Z_{\text{crit}}=2.58$

Normal Distribution

(Population Standard Deviation σ is known
or Standard Deviation σ is known but n is Large)

Normal Distribution

σ is known

$[n < 30$ (small) or $n \geq 30$ (large)]

$$z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

Normal (σ is unknown) or unknown or non-normal distribution
but $n \geq 30$ (large)

$$z = \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

← الانحراف المعياري
للعيينة

Normal Distribution

(Population Standard Deviation σ is Unknown and n is small)

Normal Distribution

σ is unknown

$N < 30$ (small)


$$t = \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

- **معرفة (σ) وحجم العينة كبير أو صغير** $\Rightarrow$ استخدام إحصائية (Z).
- **عدم معرفة (σ) أو التوزيع غير طبيعي والعينة صغيرة** $\Rightarrow$ استخدام إحصائية (t).
- **عينة كبيرة مع عدم معرفة (σ)** $\Rightarrow$ يمكن استخدام (Z) كاقتراب.

Summary of Forms for Null and Alternative Hypotheses about a Population Mean

The equality part of the hypotheses always appears in the null hypothesis H_0 . In general, a hypothesis test about the value of a population mean μ must take one of the following three forms (where μ_0 is the hypothesized value of the population mean).

a) One – tailed test

- i. $H_0: \mu = \mu_0$ vs $H_1: \mu < \mu_0$ (less than or smaller than)
- ii. Upper (right)-tailed test
 $H_0: \mu = \mu_0$ vs $H_1: \mu > \mu_0$ (more than or greater than)


b) Two – tailed test

$$H_0: \mu = \mu_0 \text{ vs } H_1: \mu \neq \mu_0 \text{ (does not equal or different from)}$$


One- vs. Two-Tailed Tests

- In theory, you should use one-tailed when ^{الاجاه المتاحه}
 1. Change in opposite direction would be meaningless
 2. Change in opposite direction would be uninteresting
 3. No opposing theory predicts change in opposite direction
- By convention/default in the social sciences, two-tailed is standard
- Why? Because it is a more strict criterion ^{لانه اختيار افضل} A more conservative test.

One tail



Two tail



Values that differ significantly from 100


Step 5

- Compare the test statistics value with the critical value of rejection.

Does Z-value = or $\neq$ value from the table

Step 6

- Decide whether to reject the null hypothesis and confidence statement.




Hypothesis Testing

Two-tailed

$$H_0: \mu = 23$$


$$H_1: \mu \neq 23$$



Left-tailed

$$H_0: \mu \geq 23$$

$$H_1: \mu < 23$$




One-tailed

Right-tailed

$$H_0: \mu \leq 23$$

$$H_1: \mu > 23$$



Rejection Rule

Lower (left)-tailed test

Hypotheses

$$H_0: \mu = \mu_0$$

vs $H_1: \mu < \mu_0$ (less than or smaller than)

Rejection Rule

Reject H_0 at a level of significance α if:

- i. $Z < -Z_{1-\alpha}$ (In case of using Z-distribution)
- ii. $t < -t_{(\alpha, n-1)}$ (In case of using t-distribution)

Rejection Rule

Upper (right)-tailed test

Hypotheses

$$H_0: \mu = \mu_0$$

vs $H_1: \mu > \mu_0$ (more than or greater than)

Rejection Rule

Reject H_0 at a level of significance α if:

- i. $Z > Z_{1-\alpha}$ (In case of using Z-distribution)
- ii. $t > t_{(\alpha, n-1)}$ (In case of using t-distribution)

Rejection Rule

Two-tailed test

Hypotheses

$$H_0: \mu = \mu_0$$

vs $H_1: \mu \neq \mu_0$ (does not equal or different from)


Rejection Rule

Reject H_0 at a level of significance α if:


i. $|Z| > Z_{1-\frac{\alpha}{2}}$ (In case of using Z-distribution)

ii. $|t| > t_{\left(\frac{\alpha}{2}, n-1\right)}$ (In case of using t-distribution)


$$H_a : \mu < \mu_0$$




$$H_a : \mu > \mu_0$$



$$H_a : \mu \neq \mu_0$$



Common Critical Values



α = tail area	central area = $1 - 2\alpha$	z_α
0.10	0.80	$z_{.10} = 1.28$
0.05	0.90	$z_{.05} = 1.645$
0.025	0.95	$z_{.025} = 1.96$
0.01	0.98	$z_{.01} = 2.33$
0.005	0.99	$z_{.005} = 2.58$

α	$1 - \alpha$	$z_{1-\alpha}$
0.10	0.90	$z_{0.90} = 1.28$
0.05	0.95	$z_{0.95} = 1.645$
0.01	0.99	$z_{0.99} = 2.33$

Example Weight

Salem believes that his “true weight” is 72kg with a standard deviation of 3kg.

Salem weighs himself once a week for four weeks. The average of these four measurements is 75.4kg.


✿ Are the data consistent with Salem’s belief?



Example Weight

1. $H_0: \mu = 72$ $H_1: \mu > 72$ This is a one tail test
2. $\alpha = 0.05$
3. $\mu > 72$ (one tail test)
4. $Z_{\text{crit}} = Z_{\text{Reject } H_0} \text{ if } z \geq 1.645$
5. $Z = \frac{\bar{X} - \mu_0}{s/\sqrt{n}} = \frac{75.4 - 72}{3/\sqrt{4}} = 2.26$ $P(Z > 2.26) = .012$
6. Since $2.26 > 1.645$, we Reject H_0 . There is a statistically significant evidence at $\alpha = 0.05$ to show that the mean weight measured is higher than his original belief about his weight. The chance that the measured weight and initial (belief) weight means are different due to chance only is less than 5%.

Example Weight illustrated



Example

Researchers are interested in the mean age of a certain population. They are wondering if the **mean age** is more than **25** years. Assuming that the **population is normally distributed** with variance equal to 20. A **random sample of 10** individuals drawn from the population of interest. From this sample, a mean of 27 is calculated. Construct the proper hypothesis, test your hypothesis, and then state the proper conclusion? Use $\alpha = 0.05$ to test the hypothesis?

Solution

We have:

1- Normal distribution.

2- The standard deviation σ is known.

Then we will use the standard normal distribution (Z).

$$n = 10, \mu_0 = 25, \bar{X} = 27, \sigma = \sqrt{\sigma^2} = \sqrt{20} = 4.472, \alpha = 0.05$$

Example

Hypotheses

$$H_0: \mu = \mu_0 \rightarrow H_0: \mu = 25$$

$$\text{vs } H_1: \mu \geq \mu_0 \rightarrow H_1: \mu > 25 \text{ (more than)}$$

Rejection Rule

Reject H_0 at a level of significance $\alpha = 0.05$ if:

$$Z > Z_{1-\alpha}, Z_{1-\alpha} = Z_{1-0.05} = Z_{0.95} = 1.645$$

Test statistics (calculated value)

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} = \frac{27 - 25}{4.472 / \sqrt{10}} = 1.414$$

6 points

Continued

Decision

We get $Z = 1.414 < Z_{1-\alpha} = 1.645$

Then the rejection is not satisfied and therefore the decision will be do not reject (accept) H_0 at $\alpha = 0.05$ and therefore we conclude that the mean age is 25 years, that is, $\mu = 25$ years

Example

The **mean** maximum oxygen uptake for a sample of **242** women was **32.3** with a **standard deviation** of **12.14**, we wish to know if, on the basis of the data, can we conclude that the **mean** score for a population of such women is **smaller** than **33**? Use **$\alpha = 0.01$** to test the hypothesis?

Solution

We have:

- 1- Unknown distribution (population).*
 - 2- The standard deviation σ is unknown ($S = 12.14$).*
 - 3- The sample size (n) is large ($n = 242 > 30$).*
- Then we will use the standard normal distribution (Z).

$$n = 242, \mu_0 = 33, \bar{X} = 32.30, S = 12.14, \alpha = 0.01$$

Continued

Lower (Left) -Tailed Test

Hypotheses

$$H_0 : \mu = 33$$

vs $H_1 : \mu < 33$ (smaller than).

Rejection Rule

Reject H_0 at level of significance $\alpha = 0.01$ if :

$$Z < -Z_{1-\alpha}$$

Test Statistic (*Calculated Value*)

$$Z = \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

$$\Rightarrow Z = \frac{32.30 - 33}{12.14 / \sqrt{242}} = -0.897$$

Decision

We get $Z = -0.897 > -Z_{1-\alpha} = -2.33$

Then the rejection rule is not satisfied and therefore the decision will be do not reject (accept) H_0 at $\alpha = 0.01$ and therefore we conclude that the mean score for a population of such women is 33, that is, $\mu = 33$.

Critical Value (*Tabulated Value*)

$$\begin{aligned} Z_{1-\alpha} &= Z_{1-0.01} = Z_{0.99} = 2.33 \\ \Rightarrow -Z_{1-\alpha} &= -Z_{1-0.01} = -Z_{0.99} = -2.33 \end{aligned}$$

Example

The body mass index (BMI) of a **group** of **14** healthy adult males has a **mean of 30.5** and a **standard deviation of 10.6392**, can we conclude that the **mean BMI** of the **population** is equal to **36** assuming that the population is normally distributed? Use $\alpha = 0.1$ to test the hypothesis?

Solution

We have:

- 1- Normal distribution (Normal population).
- 2- The standard deviation σ is unknown
($S = 10.6392$).
- 3- The sample size (n) is small ($n = 14 < 30$).

➡ Then we will use the t-distribution.

$$n = 14, \mu_0 = 36, \bar{X} = 30.5, S = 10.6392, \alpha = 0.10$$

Two -Tailed Test

Hypotheses

$$H_0 : \mu = 36$$

vs $H_1 : \mu \neq 36$ (does not equal).

Rejection Rule

Reject H_0 at level of significance $\alpha = 0.10$ if :

$$|t| > t_{(\frac{\alpha}{2}, n-1)}$$

Test Statistic (Calculated Value)

$$t = \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

$$\Rightarrow t = \frac{30.5 - 36}{10.6392 / \sqrt{14}} = -1.934$$

Critical Value (Tabulated Value)

$$t_{(\frac{\alpha}{2}, n-1)}$$

$$= t_{(\frac{0.10}{2}, 14-1)}$$

$$= t_{(0.05, 13)}$$

$$= 1.771$$

Decision

We get $|t| = |-1.934| = 1.934 > t_{(\frac{\alpha}{2}, n-1)} = t_{(0.05, 13)} = 1.771$

Then the rejection rule is satisfied and therefore the decision will be **reject** H_0 at $\alpha = 0.10$ and therefore we conclude that the mean BMI of the population is not equal to 36, that is, $\mu \neq 36$. In other words H_1 is accepted at $\alpha = 0.10$.