



PHYSIOLOGY

FACULTY OF PHARMACEUTICAL SCIENCES

DR. AMJAAD ZUHIER ALROSAN

LECTURE 9-PARTS (1) & (2): STRUCTURE AND FUNCTION OF BLOOD VESSELS AS WELL AS
FACTORS AFFECTING BLOOD FLOW AND BLOOD PRESSURE

Objectives

1. Discuss **structure and function of blood vessels.**
2. Describe **capillary exchange.**
3. Explore **hemodynamics: factors affecting blood flow.**


(Pages 730- 743 of the reference)

Objectives

4. Discuss control of blood pressure and blood flow.
5. Describe checking circulation.
6. Explore shock and homeostasis.

(Pages 744- 753 of the reference)

tunica interna
tunica media
tunica externa



STRUCTURE AND FUNCTION OF BLOOD VESSELS




Diagram illustrating the structure of a blood vessel wall. The layers from the outside in are: elastic, muscular, postcapillaries, and muscular.

- The five main types of blood vessels are **arteries**, **arterioles**, **capillaries**, **venules**, and **veins**.
- **Arteries** *carry blood away from the heart to other organs*. Large, elastic arteries leave the heart and divide into medium-sized, muscular arteries that branch out into the various regions of the body. Medium-sized arteries then divide into small arteries, which in turn divide into still smaller arteries called **arterioles**. *(smooth muscles)*
- As the **arterioles enter a tissue**, they branch into numerous tiny vessels called **capillaries**. The thin walls of capillaries allow the exchange of substances between the blood and body tissues. **Groups of capillaries within a tissue reunite to form small veins called venules**. These in turn merge to form progressively larger blood vessels called **veins**. **Veins are the blood vessels that convey blood from the tissues back to the heart.**

BASIC STRUCTURE OF A BLOOD VESSEL

The three structural layers of a generalized blood vessel **from innermost to outermost** are the tunica interna (intima), tunica media, and tunica externa (adventitia).

external elastic lamina ptiwbéig!

Kidneys doesn't have
tunica externa

TUNICA INTERNA (INTIMA)

- ❖ The **tunica interna (intima)** forms the **inner lining** of a blood vessel and is **in direct contact with the blood** as it flows **through the lumen**, or **interior opening**, of **the vessel**. Although this layer has multiple parts, these tissue components contribute minimally **to the thickness of the vessel wall**.
- ❖ Its **innermost layer is called endothelium**, is a **thin layer** of flattened cells that lines the inner surface of the entire cardiovascular system (**heart and blood vessels**). The **endothelial cells** are active participants in a variety of vessel-related activities, including **physical influences** on **blood flow**, **secretion of locally acting chemical mediators** that influence the contractile state of the vessel's overlying smooth muscle, and **assistance with capillary permeability**. In addition, their **smooth luminal surface** **facilitates efficient blood flow** by **reducing surface friction**. ★

TUNICA INTERNA (INTIMA)

- ❖ The second component of the tunica interna is a **basement membrane** deep to the endothelium. It provides a physical support base for the epithelial layer. It appears to play an important role in guiding cell movements during tissue repair of blood vessel walls. ²
- ❖ The outermost part of the tunica interna, which forms the boundary ^{geo} _{1,2,3} between the tunica interna and tunica media, is the **internal elastic lamina**, is a thin sheet of elastic fibers with a variable number of windowlike openings that facilitate diffusion of materials through the tunica interna to the thicker tunica media.

TUNICA MEDIA


- ✓ Is the tissue layer that displays the **greatest variation among the different vessel types**. In most vessels, it is a relatively thick layer comprising mainly smooth muscle cells and substantial amounts of elastic fibers.
- ✓ An increase in sympathetic stimulation typically stimulates the smooth muscle to contract, squeezing the vessel wall and narrowing the lumen. Such a decrease in the diameter of the lumen of a blood vessel is called **vasoconstriction**.
- ✓ In contrast, when sympathetic stimulation decreases, smooth muscle fibers relax. The resulting increase in lumen diameter is called **vasodilation**.

sympathetic ↑ HR ↑ contractility ↑ vasoconstriction peripherally


” ↓ HR ↓ relaxation ↑ vasodilation peripherally

TUNICA EXTERNA

↳ elastic fibers/collagen fibers



The outer covering of a blood vessel, the tunica externa, consists of **elastic and collagen fibers**.



Function :

①

In addition to the important role of supplying the vessel wall with nerves and self-vessels, the tunica externa **helps anchor the vessels to surrounding tissues**.

②

endothelium cells & 4 Function

↳ direct contact with blood

1- smooth blood flow (minimize friction)

2. maintain vascular tone  contraction (Balanced)
 relaxation

vasoconstricting + vasodilation happens in **tunica media**

(vaso motor center)

smooth muscles / elastic fibers

How ?! two ways

secrete vasoconstrictors and vasodilators by itself

regulation of BP is 3 types: 1- Auto regulation
2- short term regulation
3- long term regulation

b transocytosis

لکھنؤ

substances → endothelial cells → basement membrane → internal elastic lamina

through transcytosis (endocytosis)

↓

exocytosis

tunica media (elastic fibers)

↓

tunica externa (support)

السراويل
الحبل

↓

تحت

3 - coagulation regulation

grow coagulation factors / anti coagulation Factors

→ grow) inflammatory factors / anti inflammatory Factors


L Balanced

4- inflammation regulation


- إلى يحتمم الدكتور بالاستماعات -

- branching of blood vessels (التفرع بالإنجليزية)
downward little by little diameter \downarrow pressure -
this may be called regulation of blood pressure + blood flow regulation -
- the 3 layers of blood vessels (الطبقة)
- the 4 functions of endothelial cells


smooth muscles (in tunica media) vaso constriction (narrowing)
vasodilation



ARTERIES



Their walls stretch easily or expand without tearing in response to a small increase in pressure (due to their plentiful elastic fibers).



The wall of an artery has the three layers of a typical blood vessel.

ELASTIC ARTERIES

- The **largest arteries in the body**.
- **TUNICA INTERNA:** Well-defined internal elastic lamina.
- **TUNICA MEDIA:** Thick and dominated by elastic fibers; well-defined external elastic lamina.
- **TUNICA EXTERNA:** Thinner than tunica media.
- **Function:** **Conduct blood from heart to muscular arteries**.

MUSCULAR ARTERIES

- **The Medium-sized arteries.**
- **TUNICA INTERNA:** Well-defined internal elastic lamina.
- **TUNICA MEDIA:** Thick and dominated by smooth muscle; thin external elastic lamina.
- **TUNICA EXTERNA:** Thicker than tunica media.
- **Function:** **Distribute blood to arterioles.**

g-jg

ARTERIOLES

- **Microscopic (15–300 micrometer in diameter).**
- **TUNICA INTERNA:** Thin with a fenestrated internal elastic lamina that disappears distally.
- **TUNICA MEDIA:** One or two layers of circularly oriented smooth muscle;
الأخير distalmost smooth muscle cell forms a precapillary sphincter.
- **TUNICA EXTERNA:** Loose collagenous connective tissue and sympathetic nerves.
1 2
- **Function:** Deliver blood to capillaries and help regulate blood flow from arteries to capillaries.

CAPILLARIES

- Microscopic; smallest blood vessels (5–10 micrometer in diameter).
- TUNICA INTERNA: Endothelium and basement membrane.
- TUNICA MEDIA: None.
- TUNICA EXTERNA: None.
- Function: Permit exchange of nutrients and wastes between blood and interstitial fluid; distribute blood to postcapillary venules.
 - ① تنس
 - ② توزع

POSTCAPILLARY VENULES

- **Microscopic** (10–50 micrometer in diameter).
- **TUNICA INTERNA:** Endothelium and basement membrane.
- **TUNICA MEDIA:** None.
- **TUNICA EXTERNA:** Sparse. متفرق
- **Function:** ^①Pass blood into muscular venules; ^②permit exchange of nutrients and wastes between blood and interstitial fluid and ^③function in white blood cell emigration.

احجر

MUSCULAR VENULES

- **Microscopic** (50–200 micrometer in diameter).
- **TUNICA INTERNA:** Endothelium and basement membrane.
- **TUNICA MEDIA:** One or two layers of circularly oriented smooth muscle.
- **TUNICA EXTERNA:** Sparse. *← sanguino*
- **Function:** Pass blood into vein; act as reservoirs for accumulating large volumes of blood (along with postcapillary venules).

VEINS

- Range from 0.5 mm to 3 cm in diameter.
- **TUNICA INTERNA:** Endothelium and basement membrane; no internal elastic lamina; contain valves; lumen much larger than in accompanying artery.
- **TUNICA MEDIA:** Much thinner than in arteries; no external elastic lamina.
- **TUNICA EXTERNA:** Thickest of the three layers.
- **Function:** Return blood to heart, facilitated by valves in limb veins.

12. Which of the following materials is found in the walls of capillaries?

- A. endothelium
- B. elastic fibres
- C. collagen fibres
- D. smooth muscle

Answer is A: Only endothelium (endothelial cells and the basement membrane) compose the walls of capillaries.

CAPILLARY EXCHANGE

- Is the movement of substances between blood and interstitial fluid.
- Substances enter and leave capillaries by three basic mechanisms: **diffusion, transcytosis, and bulk flow.**
 - 1- filtration
 - 2- reabsorption

DIFFUSION

- The most important method of capillary exchange is simple diffusion.
- Because O₂ and nutrients normally are present in higher concentrations in blood, they diffuse down their concentration gradients into interstitial fluid and then into body cells. CO₂ and other wastes released by body cells are present in higher concentrations in interstitial fluid, so they diffuse into blood.

TRANSCYTOSIS

- A small quantity of material crosses capillary walls by transcytosis.
- In this process, substances in blood plasma become enclosed within tiny pinocytic vesicles that first enter endothelial cells by endocytosis, then move across the cell and exit on the other side by exocytosis.

BULK FLOW: FILTRATION AND REABSORPTION

- Bulk flow is a **passive process** in which large numbers of ions, molecules, or particles in a fluid move together in the same direction.
- Bulk flow occurs from an area of higher pressure to an area of lower pressure, and it continues as long as a pressure difference exists. ★
- Diffusion is more important for **solute exchange** between blood and interstitial fluid, but **bulk flow** is more important for **regulation of the relative volumes of blood and interstitial fluid.**
- Pressure-driven movement of fluid and solutes from blood capillaries into interstitial fluid is called **filtration**. Pressure-driven movement from interstitial fluid into blood capillaries is called **reabsorption.**

HEMODYNAMICS: FACTORS AFFECTING BLOOD FLOW

- **Blood flow** is the volume of blood that flows through any tissue in a given time period (in mL/min). **Total blood flow is cardiac output (CO)**, the volume of blood that circulates through systemic (or pulmonary) blood vessels each minute.
- How the cardiac output becomes distributed into circulatory routes that serve various body tissues depends on **two more factors**:
 - (1) the pressure difference that drives the blood flow through a tissue.
 - (2) the resistance to blood flow in specific blood vessels.

pto

- Blood flows from regions of higher pressure to regions of lower pressure; **the greater the pressure difference, the greater the blood flow. But the higher the resistance, the smaller the blood flow.**

Elmuse

BLOOD PRESSURE

- **Contraction of the ventricles generates blood pressure (BP), the hydrostatic pressure exerted by blood on the walls of a blood vessel.**
- BP is determined by cardiac output, blood volume, and vascular resistance.
- BP is highest in the aorta and large systemic arteries; in a resting, young adult, BP rises to about 110 mmHg during systole (ventricular contraction) and drops to about 70 mmHg during diastole (ventricular relaxation).
- **Mean arterial pressure (MAP), the average blood pressure in arteries, is roughly one-third of the way between the diastolic and systolic pressures. It can be estimated as follows:**

- pulse pressure -

$$\text{MAP} = \text{diastolic BP} + \frac{1}{3} (\text{systolic BP} - \text{diastolic BP})$$

BLOOD PRESSURE

- We have already seen that cardiac output equals heart rate multiplied by stroke volume. Another way to calculate cardiac output is to divide mean arterial pressure (MAP) by resistance (R): $CO = \frac{MAP}{R}$. By rearranging the terms of this equation, you can see that MAP = CO * R.
- Blood pressure also depends on the total volume of blood in the cardiovascular system. The normal volume of blood in an adult is about 5 liters. Any decrease in this volume, as from hemorrhage, decreases the amount of blood that is circulated through the arteries each minute.
نزيف
- A modest decrease can be compensated for by homeostatic mechanisms that help maintain blood pressure, but if the decrease in blood volume is greater than 10% of the total, blood pressure drops. Conversely, anything that increases blood volume, such as water retention in the body, tends to increase blood pressure.

بُطَاد

water retention

BP ↑

VASCULAR RESISTANCE

- Vascular resistance is the opposition to blood flow due to friction between blood and the walls of blood vessels.
- Vascular resistance depends on (1) **size of the blood vessel lumen**, (2) **blood viscosity**, and (3) **total blood vessel length**.

SIZE OF THE LUMEN

تجويفاً

- The smaller the lumen of a blood vessel, the greater its resistance to blood flow.
- Resistance is inversely proportional to the fourth power of the diameter (d) of the blood vessel's lumen.

BLOOD VISCOSITY

- The viscosity of blood depends mostly on the ratio of red blood cells to plasma (fluid) volume, and to a smaller extent on the concentration of proteins in plasma.
- The higher the blood's viscosity, the higher the resistance. حجم خلايا الدم
- Any condition that increases the viscosity of blood, such as dehydration or polycythemia (an unusually high number of red blood cells), thus increases blood pressure.
- A depletion of plasma proteins or red blood cells, due to anemia or hemorrhage, decreases viscosity and thus decreases blood pressure.

نها

SYSTEMIC VASCULAR RESISTANCE (SVR)

- Resistance to blood flow through a vessel is directly proportional to the length of the blood vessel.
- The longer a blood vessel, the greater the resistance.
- Systemic vascular resistance (SVR), also known as total peripheral resistance (TPR), refers to all of the vascular resistances offered by systemic blood vessels.
- The diameters of arteries and veins are large, so their resistance is very small because most of the blood does not come into physical contact with the walls of the blood vessel. The smallest vessels—arterioles, capillaries, and venules—contribute the most resistance.


Determinants of Resistance:

Blood Viscosity (η) \propto Resistance



مقدار η \rightarrow مقدار \downarrow

Vessel Length (l) \propto Resistance



مقدار l \rightarrow مقدار \uparrow

Vessel Radius (r) \propto Resistance



مقدار r \rightarrow مقدار \uparrow


Factors Affecting Blood Flow

①

Pressure and resistance both affect blood flow to tissues, but they have opposing effects.

Blood flow and pressure are directly related: when pressure increases, flow increases.


Blood flow and resistance are inversely related: when resistance increases, flow decreases.



VELOCITY OF BLOOD FLOW

- Earlier we saw that blood flow is the volume of blood that flows through any tissue in a given time period (in mL/min).
- The speed or velocity of blood flow (in cm/sec) is inversely related to the cross-sectional area.
- **Velocity is slowest where the total cross-sectional area is greatest.**
- Each time an artery branches, the total cross-sectional area of all of its branches is greater than the cross-sectional area of the original vessel, **so blood flow becomes slower and slower as blood moves further away from the heart, and is slowest in the capillaries.**

Factors that Increase Blood Pressure



Physiological factors affecting blood pressure

Factor	Effect on blood pressure
The pumping action of the heart	The greater the cardiac output, the higher the arterial pressure
The blood volume	The greater the blood volume, the higher the arterial pressure
The viscosity of the blood	The more viscous the blood, the higher the arterial pressure
The condition of the blood vessels (resistance)	The greater the resistance, the higher the arterial pressure

CONTROL OF BLOOD PRESSURE AND BLOOD FLOW

- Several interconnected **negative feedback systems** control blood pressure by adjusting heart rate, stroke volume, systemic vascular resistance, and blood volume.
- Some systems allow **rapid adjustments** to cope with sudden changes, such as the drop in blood pressure in the brain that occurs when you get out of bed; others act more slowly to provide long-term regulation of blood pressure.

CONTROL OF BLOOD PRESSURE AND BLOOD FLOW

□ Role of the Cardiovascular Center in the medulla oblongata:

- ❖ 1- helps regulate heart rate and stroke volume.
- ❖ 2- controls neural, hormonal, and local negative feedback systems that regulate blood pressure and blood flow to specific tissues.
- ❖ Groups of neurons scattered within the CV center:
 - ❖ Some neurons stimulate the heart (cardiostimulatory center); others inhibit the heart (cardioinhibitory center).
 - ❖ Still others control blood vessel diameter by causing constriction (vasoconstrictor center) or dilation (vasodilator center).

1. ROLE OF THE CARDIOVASCULAR CENTER

- ❖ The cardiovascular center receives input both **from higher brain regions and from sensory receptors**. Nerve impulses descend from the cerebral cortex, limbic system, and hypothalamus to affect the cardiovascular center.
- ❖ The three main types of sensory receptors that provide input to the cardiovascular center are **proprioceptors, baroreceptors, and chemoreceptors**.
- ❖ **Proprioceptors** monitor **movements of joints and muscles** and **provide input to the cardiovascular center during physical activity**.
- ❖ **Baroreceptors** monitor **changes in pressure and stretch in the walls of blood vessels**.
- ❖ **Chemoreceptors** monitor **the concentration of various chemicals** in the blood.

CV Center




Figure 21.12 Tortora - PAP 12/e
Copyright © John Wiley and Sons, Inc. All rights reserved.

ROLE OF THE CARDIOVASCULAR CENTER

- ❖ Output from the cardiovascular center flows along sympathetic and parasympathetic neurons of the ANS.
- ✓ Sympathetic impulses reach the heart via the cardiac accelerator nerves. An increase in sympathetic stimulation increases heart rate and contractility.
 $HR \uparrow$
- ✓ Parasympathetic stimulation, conveyed along the vagus (X) nerves, decreases heart rate.
 $HR \downarrow$

ROLE OF THE CARDIOVASCULAR CENTER


- ❖ The cardiovascular center also continually sends impulses **to smooth muscle in blood vessel walls via vasomotor nerves**.
- ✓ Impulses propagate along sympathetic neurons that innervate blood vessels **in viscera and peripheral areas**.
- ✓ The result is a moderate state of tonic contraction or vasoconstriction, called **vasomotor tone**, that **sets the resting level of systemic vascular resistance**. Sympathetic stimulation of most veins causes constriction that moves blood **out of venous blood reservoirs** and increases blood pressure.

BP ↑


2. NEURAL REGULATION OF BLOOD PRESSURE

The nervous system regulates blood pressure via negative feedback loops that occur as two types of reflexes: baroreceptor reflexes and chemoreceptor reflexes.

BARORECEPTOR REFLEXES




Baroreceptors are **pressure-sensitive sensory receptors**. They are **located in the aorta, internal carotid arteries** (arteries in the neck that supply blood to the brain), and other large arteries in the neck and chest.



They **send impulses to the cardiovascular center to help regulate blood pressure**.


Innervation of the Heart

↳ supplying nerves to a body part




- Slow it down with **parasympathetic stimulation (X)**
- **Sensory information from baroreceptors (IX)**


vagus
glossopharyngeal



BARORECEPTOR REFLEXES

- Baroreceptors are pressure-sensitive neurons that monitor stretching.
- When blood pressure falls, the baroreceptors are stretched less, and they send nerve impulses at a slower rate to the cardiovascular center. In response, the CV center decreases parasympathetic stimulation of the heart by way of motor axons of the vagus nerves and increases sympathetic stimulation of the heart via cardiac accelerator nerves.
- Moving from a prone (lying down) to an erect position decreases blood pressure and blood flow in the head and upper part of the body. The baroreceptor reflexes, however, quickly counteract the drop in pressure. Sometimes these reflexes operate more slowly than normal, especially in the elderly.





CHEMORECEPTOR REFLEXES


- Chemoreceptors, sensory receptors that **monitor the chemical composition of blood**, are **located close to the baroreceptors** of the carotid sinus and arch of the aorta in small structures called **carotid bodies and aortic bodies**, respectively.

- These chemoreceptors **detect changes in blood level of O₂, CO₂, and H⁺**. Hypoxia (lowered O₂ availability), acidosis (an increase in H⁺ concentration), or hypercapnia (excess CO₂) **stimulates the chemoreceptors to send impulses to the cardiovascular center**. In response, the CV center **increases sympathetic stimulation to arterioles and veins**, producing **vasoconstriction** and an **increase in blood pressure**. These chemoreceptors also **provide input to the respiratory center** in the brain stem to adjust the rate of breathing.

sympathetic ↑ BP ↑ vasoconstriction
peripherally

Chemoreceptor

- Chemosensitive cells that respond to changes in pCO_2 and pO_2 and pH levels (Hydrogen ion).



bloodflow to Kidneys ↓ → renin \rightarrow goes يغزو ① RAAS

by juxtaglomerular cells

bloodflow to liver ↓ → angiotensin \rightarrow goes يغزو

Angiotensin I \rightarrow goes يحول (to) renin

lungs \leftarrow Angiotensin-converting enzyme \rightarrow goes to lungs

convert Angiotension I to Angiotension II

reabsorption \leftarrow Aldosterone \rightarrow goes يغزو

water / sodium ↓

distal convoluted tubule \rightarrow goes

collecting duct ↓

capillaries \rightarrow goes

renal vein \rightarrow posterior vena cava (right atrium)

method ② in regulating $BP \uparrow$

proximal convoluted tubule: Ca^{2+} reabsorption due to Angiotensin II

tubule

③ sympathetic NS: centrally dilation

peripherally constriction (Angiotensin II)

④ minimize vasodilators hormones (Atrial natriuretic peptide)

vasopressin ADH (anti diuretic hormone) → Function

Reabsorption (water + sodium)

$BF \downarrow$ \rightarrow (thirst centers) سعفہ

osmoreceptors in hypothalamus → posterior pituitary (action potential) دفعہ

(ADH) دفعہ ← gland

reabsorption through distal convoluted tubule + collecting duct

BF BP ↑

(Hypertension)

بجز عروق لينز الatrium (right atrium) (atrial natriuretic peptide)

Blood volume ↓ → Angiotensin II activation → Aldosterone activation → EP/NOR EP activation

antidiuretic hormone activation \rightarrow atrial natriuretic peptide Inhibition

Blood volume ↑ → اعجم الـ موـقـع

BP ↑ hypertension ادویه لعلی

- angiotensin-converting enzyme inhibitors
- diuresis
- angiotensin II inhibitors (antagonist)
- Beta blockers

3- HORMONAL REGULATION OF BLOOD PRESSURE

1- Renin–angiotensin–aldosterone (RAA) system: → long term regulation

BF

- When blood volume falls or blood flow to the kidneys decreases, **juxtaglomerular cells in the kidneys secrete renin into the bloodstream.**
- In sequence, ^{Kidney} renin and ^{lungs} angiotensin-converting enzyme (ACE) act on their substrates to produce the active hormone **angiotensin II**, which **raises blood pressure in two ways.**
- First**, angiotensin II is a **potent vasoconstrictor**; it raises blood pressure by increasing **systemic vascular resistance.**
- Second**, it **stimulates secretion of aldosterone**, which **increases reabsorption of sodium ions and water by the kidneys**. The water reabsorption increases **total blood volume**, which increases blood pressure.

3- HORMONAL REGULATION OF BLOOD PRESSURE

2- Epinephrine and norepinephrine:

- In response to sympathetic stimulation, the adrenal medulla releases epinephrine and norepinephrine.
- These hormones **increase cardiac output** by **increasing the rate and force of heart contractions**.

(vasopressin) contractility ↑ CO ↑ EP/NOREP ↑

3- Antidiuretic hormone (ADH):

- It is produced by the hypothalamus and released from the posterior pituitary in response to dehydration or decreased blood volume.
- It causes in an **increase in blood volume and a decrease in urine output**.

3- HORMONAL REGULATION OF BLOOD PRESSURE

4- Atrial natriuretic peptide (ANP):

- Released by cells in the atria of the heart, atrial natriuretic peptide (ANP) lowers blood pressure by causing vasodilation and by promoting the loss of salt and water in the urine, which reduces blood volume.

Function : 1- lowered BP ↓ vasodilation

2- reduce BV ↓ lose of salt + water (urine)

4- AUTOREGULATION OF BLOOD FLOW

- In each capillary bed, local changes can regulate vasomotion.
- When **vasodilators** produce local dilation of arterioles and relaxation of precapillary sphincters, blood flow into capillary networks is increased, which increases O₂ level.
$$BF \uparrow \quad O_2 \uparrow$$
- Vasoconstrictors have the opposite effect.
$$BF \downarrow \quad O_2 \downarrow$$
- **The ability of a tissue to automatically adjust its blood flow to match its metabolic demands is called autoregulation.**

4- AUTOREGULATION OF BLOOD FLOW

- **Autoregulation** also controls regional blood flow in the brain; blood distribution to various parts of the brain changes dramatically for different mental and physical activities.
- During a conversation, for example, blood flow increases to your motor speech areas when you are talking and increases to the auditory areas when you are listening.

4- AUTOREGULATION OF BLOOD FLOW

- Two general types of stimuli cause autoregulatory changes in blood flow:

1. **Physical changes.** Warming promotes vasodilation, and cooling causes vasoconstriction. In addition, smooth muscle in arteriole walls exhibits a myogenic response— it contracts more forcefully when it is stretched and relaxes when stretching lessens.
Glue
2. **Vasodilating and vasoconstricting chemicals.** Several types of cells—including white blood cells, platelets, smooth muscle fibers, macrophages, and endothelial cells—release a wide variety of chemicals that alter blood-vessel diameter.

4- AUTOREGULATION OF BLOOD FLOW

- An important difference between the pulmonary and systemic circulations is their autoregulatory response to changes in O₂ level.

قریب

- The walls of blood vessels in the systemic circulation dilate in response to low O₂. With vasodilation, O₂ delivery increases, which restores the normal O₂ level.

centrally

بعد عن القلب

peripherally

- By contrast, the walls of blood vessels in the pulmonary circulation constrict in response to low levels of O₂. This response ensures that blood mostly bypasses those alveoli (air sacs) in the lungs that are poorly ventilated by fresh air. Thus, most blood flows to better-ventilated areas of the lung.

زجا

CHECKING CIRCULATION

1. Pulse:

- The alternate expansion and recoil of elastic arteries after each ^{contraction} systole of the left ventricle creates a traveling pressure wave that is called the pulse.
- The pulse is strongest in the arteries closest to the heart, becomes weaker in the arterioles, and disappears altogether in the capillaries.
- The pulse rate normally is the same as the heart rate, about 70 to 80 beats per minute at rest. **Tachycardia** is a rapid resting heart or pulse rate over 100 beats/min. **Bradycardia** is a slow resting heart or pulse rate under 50 beats/min.

CHECKING CIRCULATION

2. Measuring Blood Pressure:

- The term **blood pressure** usually refers to the pressure in arteries generated by the left ventricle during systole and the pressure remaining in the arteries when the ventricle is in diastole. Blood pressure is usually measured in the brachial artery in the left arm.
- The device used to measure blood pressure is a **sphygmomanometer**.
- The normal blood pressure of an adult male is less than 120 mmHg systolic and less than 80 mmHg diastolic. For example, “110 over 70” (written as 110/70) is a normal blood pressure.

CHECKING CIRCULATION

2. Measuring Blood Pressure:

$$(\text{systolic BP} - \text{diastolic BP}) = \text{pulse}$$

- The difference between systolic and diastolic pressure is called **pulse pressure**. This pressure, normally about **40 mmHg**, provides information about the condition of the **cardiovascular system**. For example, conditions such as **atherosclerosis** greatly increase **pulse pressure**. The normal ratio of systolic pressure to diastolic pressure to pulse pressure is about 3:2:1.

SHOCK AND HOMEOSTASIS

- ❑ Shock is a failure of the cardiovascular system to deliver enough O₂ and nutrients to meet cellular metabolic needs.
- ❑ The causes of shock are many and varied, but all are characterized by inadequate blood flow to body tissues. With inadequate oxygen delivery, cells switch from aerobic to anaerobic production of ATP, and lactic acid accumulates in body fluids. If shock persists, cells and organs become damaged, and cells may die unless proper treatment begins quickly.

TYPES OF SHOCK

➤ Shock can be of four different types:

- (1) **Hypovolemic shock** due to decreased blood volume
- (2) **Cardiogenic shock** due to poor heart function.
- (3) **Vascular shock** due to inappropriate vasodilation.
- (4) **Obstructive shock** due to obstruction of blood flow.

انسداد



دکورہ
حذفیں

TYPES OF SHOCK

Cause	Pathophysiology*	Patterns of abnormalities		
		Filling status	Cardiac function	Systemic resistance
Hypovolemic	Loss of volume	low	low	high
Vasogenic	Vasodilation	low	high	low
Cardiogenic	Pump failure	high	low	high
Obstructive	Obstruction to flow	Variable**	low	high

*primary problem mentioned in BOLD ; **depending on site of obstruction

HYPOVOLEMIC SHOCK

- ✓ It is due to decreased blood volume (i.e. **blood loss** as acute (sudden) hemorrhage and loss of body fluids i.e. through excessive sweating).
- ✓ Whatever the cause, when the volume of body fluids falls, venous return to the heart declines, filling of the heart lessens, stroke volume decreases, and cardiac output decreases. Replacing fluid volume as quickly as possible is essential in managing hypovolemic shock.

$\beta F \downarrow$ Filling \downarrow $SV \downarrow$ $CO \downarrow$

CARDIOGENIC SHOCK

- ① ✓ The heart fails to pump adequately, most often because of a myocardial infarction (heart attack).
- 2 - نفق دم -
- ✓ Other causes of cardiogenic shock include poor perfusion of the heart (ischemia),³ heart valve problems,⁴ excessive preload or afterload, impaired contractility of heart muscle fibers, and arrhythmias.⁵

6 عدم انتظام
حرباً ←
القلب ←

VASCULAR SHOCK

- ✓ A variety of conditions can cause inappropriate dilation of arterioles or venules.

↳

- ✓ In **anaphylactic shock**, a severe allergic reaction—for example, to a bee sting—releases histamine and other mediators that cause vasodilation.
- ✓ In **neurogenic shock**, vasodilation may occur following trauma to the head that causes malfunction of the cardiovascular center in the medulla.
- ✓ Shock stemming from certain bacterial toxins that produce vasodilation is termed **septic shock.**

OBSTRUCTIVE SHOCK

- ✓ It occurs when **blood flow through a portion of the circulation is blocked.**
→ انسداد
- ✓ The most common cause is **pulmonary embolism**, a blood clot lodged in a blood vessel of the lungs.

HOMEOSTATIC RESPONSES TO SHOCK

- The major mechanisms of **compensation in shock** are **negative feedback systems** that work to return cardiac output and arterial blood pressure to normal.
- ❖ **Activation of the renin–angiotensin–aldosterone system:** **angiotensin II** causes ¹ vasoconstriction and ² stimulates the adrenal cortex to secrete aldosterone, a hormone that increases reabsorption of sodium ions and water by the kidneys.
- ❖ **Secretion of antidiuretic hormone:** the posterior pituitary releases more antidiuretic hormone (ADH). ADH enhances water reabsorption by the kidneys, which conserves remaining blood volume.

HOMEOSTATIC RESPONSES TO SHOCK

- ❖ **Activation of the sympathetic division of the ANS:** aortic and carotid baroreceptors initiate powerful sympathetic responses throughout the body. One result is marked vasoconstriction of arterioles and veins of the skin, kidneys, and other abdominal viscera. (Vasoconstriction does not occur in the brain or heart.) Constriction of arterioles increases systemic vascular resistance, and constriction of veins increases venous return.
*arterioles SVR ↑
veins VR ↑*
- ❖ **Release of local vasodilators:** In response to hypoxia, cells liberate vasodilators—including potassium ions, hydrogen ions, lactic acid, adenosine, and nitric oxide—that dilate arterioles and relax precapillary sphincters. Such vasodilation increases local blood flow and may restore O_2 level to normal in part of the body.

HOMEOSTATIC RESPONSES TO SHOCK

- ❖ If blood volume drops more than 10–20%, or if the heart cannot bring blood pressure up sufficiently, compensatory mechanisms may fail to maintain adequate blood flow to tissues. At this point, shock becomes life-threatening as damaged cells start to die.