


مع سنوات !!



Crude drug
تخزين العقاقير الخام من النباتات الطبية

Pharmacognosy

PREPARATION OF CRUDE DRUGS FROM MEDICINAL PLANTS

Cultivation, collection and processing of
herbal drug

References

Trease and Evans, Pharmacognosy

Dr. Jamila Dibas

Dr. Talal Abu rojai

GRAPHICAL OUTLINE



By Erhirre Earnest O.

Preparation of drugs for commercial market

- Crude drugs: undergo only drying and preparation by grinding.
- Preparation of drugs for commercial market includes:
 1. Collection
 2. Harvesting الحصاد
 3. Drying (By airflow or complete drying operation)
 4. Garbling التفريغ
 5. Storage and preservation

Collection of medicinal plants

Time of collection affects both quality and quantity of active constituents.

It has been found that active constituents in plants vary in amount and nature throughout the year so

Seasonal Collection is required.

For example: Rhubarb contains no anthraquinone in winter, instead it contains anthranols, In summer, anthranols are oxidized to anthraquinones (**active constituent**) so Rhubarb collection must be in summer.



• لكل بيئة موسم حصاد معين لأنه كمية المادة الفعالة
داخل كل بيئة يختلف من وقت لآخر
مثلاً هناك لزوم تقتر موسم حصاد حيث
تعمل مع الجو كمية معينة من المادة الفعالة المطلوبة



Collection of medicinal plants

- **2. Time of the Day**
- • Flowers : Morning
- • Leaves (exceptions are available) : afternoon



- **3. Stage of maturity and age:**
- - Clove: buds
- - leaves: at flowering stage.

سنة الإثمار



Collection of plant materials:

- Suitable time for collection the amount of a constituent is usually not constant throughout the life of a plant.
- [?] Roots and rhizomes should be collected in the raining season. In most cases they must be washed free of adhering soil and sand.
- [?] Bark should be collected in the dry season.
- [?] Leaves should be collected at the flowering stage.
- [?] Flowers should be collected when fully developed and also in the morning after the morning dew had evaporated. [?] Fruits and seeds should be collected when fully ripe.
- [?] Whole plant should be collected during flowering stage

during flowering stage ← whole plants, leaves, seeds

Q4. Which of the following statements regarding the collection of medicinal plants is TRUE?

- A. Bark is collected during the rainy season, and flowers are collected before they develop.
- B. Roots and rhizomes are collected during the flowering season, while leaves are collected in the dry season.
- C. Fruits are collected before they ripen, and seeds are collected while still immature.
- D. Roots and rhizomes are collected during the dry season, while leaves are collected at the flowering stage.
- E. Whole plants are always collected after the fruits have matured.

D



يؤثر على كمية المادة التي نحصل عليها من النبات وصحة المواد النباتية
لذلك يجب أن تكون جودة المواد النباتية التي نحصل عليها عالية
الجودة أي أن تكون quantity
والكمية أي أن تكون quality

بشكل عام، عندما يصل الجزء النباتي إلى أفضل مرحلة من مراحل التطور والنمو، يُفترض أن المادة الفعالة المطلوبة تكون في أعلى تركيز لها

- Generally, when, plant part reaches it's optimal state of development, it is assumed that the targeted active constituent is in highest concentration:
- 1. **Roots and rhizomes collected at the end of the **vegetation** period (in autumn).**
- 2. **Bark is collected in the spring.** (dry season)
- 3. Leaves and herbs are collected when photosynthesis is most active and before fruit maturation.
- 4. **Flowers are usually gathered when fully developed.**
- 5. **Fruits and seeds are collected when fully-grown but unripe.**
- 6. **Seeds: fully matured and before the fruits have opened**

في نهاية موسم
النمو النباتي يعني
بفضل الخريف

2-Preservation

- Preservation aims at limiting decomposition processes as much as possible.
- The plant material must first be preserved so that the active compounds will remain unchanged during transport and storage.
- Factors that might affect the quality of crude drugs:
 - 1. Light
 - 2. Oxygen
 - 3. Attacks by insects (this can be achieved by using methyl bromide or heating up to 65C)
 - 4. Low Temperature

Q29. Which of the following does NOT affect the quality of crude drugs during storage?

- A. Oxygen
- B. Temperature
- C. Moisture
- D. Light
- E. None of the above

E

Q15. Which of the following is NOT a correct condition for the storage of crude drugs?

- A. Store under dry conditions.
- B. Protect from insects.
- C. Exclude light.
- D. Allow free exposure to air.
- E. Store in suitable containers.

0

Preservation of plant materials

- After collection, the plant material must first be preserved **so that the active compounds** will remain unchanged during transport and storage.
- The most common method for preserving plant material is **drying**. Living plant materials have high water content:
 - [?] Leaves may contain 60-90%
 - [?] Roots and rhizomes 70-85%
 - [?] Wood 40-50%
 - [?] Seed contains 5-10%

2-preservative

الدواء هو limiting decomposition التخليق من مخزون صودت تلفه الى drug
وبالتالي، يحافظ على سلامة المادة النقا له انتهاء النقل والتخزين

3- Drying of crude drugs

- **Aim of drying:**
- 1. Remove moisture since it favors spoilage due to microbial growth
- 2. Prevents molding يمنع تكون العفن
- 3. Prevents enzymatic action
- 4. Prevents bacterial action.
- 5. Prevents chemical changes
- 6. Fixes cost. يثبت التكلفة
- 7. To facilitate their grinding. يسهل عملية الطحن
- 8. To reduce their size and weight.

Drying medicinal plants?


- 1. Drying in open air: in sun and under cover sheds at night or during wet weather, e.g. clove, cinnamon,...
- 2. Artificial oven drying (Oven-drying is more rapid than air- drying, controlled temp. In oven – drying , flowers and leaves are usually dried below 60° C to retain color , while roots and barks may be dried at slightly higher temperature.
- 3. Vacuum drying: in oven, rapid and at low temp.
- 4. Lyophilization (Freeze Drying) used for biological fluids, enzymes, proteins and royal jelly. No heat involved.

Q26. Which of the following statements about the drying of crude drugs is TRUE?

- A. Roots and rhizomes should always be dried above 100°C.
- B. Leaves and flowers should be dried below 60°C to retain their color.
- C. All crude drugs are dried under direct sunlight regardless of their constituents.
- D. Fruits should never be dried after harvesting.
- E. Drying is performed only after storage.

Chilis drying





- **In order to keep crude drugs for longer periods:**
- 1. It is essential to store them in a dry condition in carefully closed containers.
- 2. It is also advisable to exclude light, because - even if it does not affect the active constituents - it almost always causes changes in the appearance of the drug, especially loss of color.
- 3. It is also necessary to protect the drug against insect attack.

4- Grinding (size reduction of plant materials):

- the first operation that must be performed after drying is grinding of the plant material to a powder of suitable particle size.
- [?] Large particles take a longer time for complete extraction than small ones and large differences in particle size thus slow down the extraction process.
- [?] The purpose for powdering the plant material is to rupture its tissue and cell structures so that its medicinal ingredients are exposed to the extraction solvent.
- [?] Furthermore, size reduction maximizes the surface area, which in turn enhances the mass transfer of active principle from plant material to the solvent.

Q16. Which of the following statements about grinding of crude drugs is TRUE?

- A. Grinding is performed only after extraction.
- B. Grinding is used to maximize particle size.
- C. Grinding minimizes the surface area of the drug.
- D. Grinding is used to maximize surface area.
- E. Grinding is carried out only for liquid drugs.

Preparation: Grinding of crude drugs

First step in preparation is grinding of the plant material to a powder of suitable particle size.

Grinding decreases the particle size and increase the surface area.

1. Mortar and pestle.

2. Hammer mill a common type for grinding crude drugs.

3. Knife mill is useful for production of low-dust powders of leaves, barks and roots for subsequent percolation or maceration.

الاستخلاص بالرشح المتكرر

النقح



➤ Machines are available for grinding crude drugs



Mortar and pestle



Hammer mill, Knife mill



Electrical blender

5- Sieving and Garbling

- Simply it is removing of external matter
- In garbling the material is passed through a sieve of suitable mesh size giving two fractions. The remaining fraction consists of coarser particles which are returned to the mill for continued grinding.



Storage

- Well-dried and well-stored plant materials can be used up to 6 months after the collection.
- Discard the plant materials if there are molds or other signs of decay. *علامات التلف*
- Use glass, ceramic pot as cooking utensils, stainless steel pot can also be used.
- Plastic may be used but not for storing crude drugs with volatile oil.
- Do not use pots made of aluminum, iron, tin or other metals as these will leach into the tea. *لا ينبغي استخدام أواني تتفاعل وتتلف الأدوية فيها*
- Use pure water (distilled water).

Q31. Which of the following statements about the storage of crude drugs is WRONG?

- A. Volatile oils should be stored in plastic containers.
- B. Crude drugs should be protected from light.
- C. Crude drugs should be stored under dry conditions.
- D. Storage containers should protect the drugs from air and moisture.
- E. Crude drugs should be protected from insect infestation.

Extraction and Separation of natural drugs

➤ INTRODUCTION

الاستخلاص

Extraction: This is the separation of medicinally active portions of plant from the inactive components by using selective solvents in standard extraction procedures.


بالاستخدام مذيبات انتقائية

Extract: This is a preparation of crude drug which contains all the constituents which are soluble in the solvent used in making the extract.

هو مستحضر يتم الحصول عليه من العقار الخام (Crude drug)، ويحتوي على جميع المكونات التي تذوب في المذيب المستخدم أثناء عملية الاستخلاص.

➤ INTRODUCTION


From Plant to Drug



➤ BASIC PRINCIPLE IN EXTRACTION

Plant constituents are usually contained inside the cells.

Therefore, The solvent used for extraction must diffuse into the cell to dissolve the desired compounds.



Extraction and Separation of natural drugs

- The commonly employed technique for removal of the active substance from the crude drug is called **extraction**
- The forms in which plant drugs are used:
 1. Fresh plant material
 2. Dried plant material
 3. Acellular products like Volatile oil, Resins, and fixed oils
 4. Galenical preparation and extracts
 5. Processed extracts

علاج طبيعي

بني منقبات ما تحتوي
على خلايا.



1. Fresh plant material

- Predominantly used in folk and herbal remedies. الطب الشعبي الطبخ بالأعشاب
- Fresh plant material is often required by the flavor industry and is also used by the perfumery industry where the volatile oils from fresh materials , such as flower petals, are highly prized and highly priced.


2. Dried plant material

The method of drying depends on the nature of the plant material and the active constituents. Plants containing volatile oils lose much of their aroma if dried at elevated temperature and, if required for their volatile content, must be dried extremely carefully. دائماً

Plant material obtained in the tropics is normally sun-dried, often being bleached during the drying process. او بيرقيوم

النباتات الاستوائية

بوتان



Processed extraction

- If the active constituents are not present in a free form but need liberating by enzymatic processes, care must be taken not to dry the plant too quickly. For example, vanilla pod, قشور الفانيليا, the source of natural vanillin, contain the phenolic glycosides glucovanillic alcohol.
اشد بحية المعالجة During the curing process, which involve slow drying, the glycoside is enzymatically hydrolised and oxidized to vanillin.


glycoside $\xrightarrow{\text{hydrolyzed/oxidized}}$ Vanillin.

Vanilla *Vanilla planifolia*



Some active constituents need to be processed (slow drying is allowed)

Liberation of active vanillin by enzymatic process from *Gentiana Luten* and Vanilla pod



3. Acellular products

They are plant isolates like volatile oil, balsams, resins and fixed oils.
مواد معزولة

Methods to extract acellular products (Volatile oils):

- 1. destructive distillation: wood of Scotch pine (used for dermititis)
- 2. **Steam distillation using Clavenger tube**: plant material is heated with water and steam is passed through the mixture and condensed. This needs a round-bottomed flask دورق that contains water or alcohol (usually water), the plant is inserted inside it with the water, then it's attached into a condenser, also a collecting tube should be there to collect the extracted oil, distillation also requires a heat source, if the treated material is very sensitive for heat (thermo labile) the steam distillation can't be used then, but other methods like:
- ~ **Hydrosteam distillation**: here the plant is separated from the water by a membrane, the plant part is inserted to the membrane and by water heating, the vapor passes this membrane to the plant material, the v.o. is then extracted.

في هذه الطريقة يتم فصل النبات عن الماء بواسطة غشاء (Membrane).

يوضع الجزء النباتي داخل الغشاء، وعند تسخين الماء:

* يمر البخار عبر الغشاء.

* يصل إلى المادة النباتية.

* يحمل معه الزيت الطيار.

* ثم يتم استخراج الزيت.

3. Acelleular products

كل ثقب فيها متصل بفتحة تنتهي إلى وعاء تجميع (Collector)، وعند دوران الأسطوانة يتم الضغط على الثمار وتحرير الزيت ليجمع

- **3. Expression** **العصر** used **pharmaceutically for citrus oils only**, fruits are turned by a stainless steel perforated roll , each pore is attached into an opening ending into a collector, upon turning these pores are induced.
- **4. Solvent extraction , using alcohol , not used in pharmaceutical industry.** طريقة غير شائعة جداً، حيث إن بعض الزيوت الطيارة تحتوي على جزء سكري، لذلك نستخدم هذه الطريقة لاستخلاص الجزء غير السكري
- **5. Enzymatic hydrolysis** : very uncommon , some **V.O. that has a sugar part** force to use this method to extract the non-sugar part (previously mentioned example: Vanilla).
- **6. Enfleurage** : in expensive **perfume industry** , an old method that uses sheets of waxes or fatty solid materials , the plant is inserted into these glassy sheets , then put in woody frames for a week or 2 weeks in a dark area , then the plant part used is removed and the sheets then are treated with alcohol to produce a concentrate of these v.o.

يستخدم في صناعة العطور الفاخرة، وهي طريقة قديمة تستخدم صفائح من الشمع أو مواد دهنية صلبة.


يتم وضع النبات داخل هذه الصفائح الزجاجية/الدهنية، ثم توضع في إطارات خشبية لمدة أسبوع أو أسبوعين في مكان مظلم.

بعد ذلك يتم إزالة الجزء النباتي، وتعالج الصفائح بالكحول لإنتاج مركز من الزيوت الطيارة.



3. Acellular products

VOLATILE OIL STEAM DISTILLATION




Clevenger tube

Figure 1. Clevenger-type apparatus — *Appareil de type Clevenger.*

3. Acellular products

VOLATILE OIL STEAM DISTILLATION

Hydro-steam distillation



Clavenger tube




Figure 1. Clavenger-type apparatus — *Appareil de type Clavenger.*

عملية التقطير لتحضير الزيوت الطيارة

VOLATILE OIL STEAM DISTILLATION

- Principle: When water is mixed with another immiscible liquid and both are boiled together boiling will occur at lower temperature than the water's boiling point
- Plant material is heated with water and steam is passed through the mixture and condensed
- Disadvantage of steam distillation:
 - 1. unstable material may be destroyed
 - 2. other substances in the plant may affect the composition of the oil
- Ex. In Citrus spp. The acidic nature cause a number of acid catalyzed rearrangements so another method should be used.

4. Galenical preparations and extracts

- Galenical: مستحضر طبي من اصل نباتي
- Galenical preparations: Are extracts of plant materials used in therapy.
- Methods of extraction to any plant:
- **A. Infusion** : ^{زقع المادة} drug stood in hot or cold water or other solvent for a short period of time.
- Ex. Preparation of plants containing volatile oil.

Methods of extraction to any plant:

الخالي

- **B. Decoction**: drug boiled with water or solvent.

خالي العقار النباتي مع الماء او مع مذيب آني

- Ex.: roots, rhizomes and barks

نقع ب س فترة طويلة

- **C. Maceration**: prolonged infusion (2-3 weeks), often in aqueous alcohol

عملية تقع بس بسحب ال solvent من المحلول

- **D. Percolation**: Maceration followed by a slow flow of fresh solvent through the ground material.


في Maceration ثم جود بطي
لمذيب جديد خلال المادة النباتية
اعطوية

- **E. Hot extraction**: repeated infusion with hot solvent example : Soxhlet extraction

تكرار عملية التقع باستخدام مذيب ساخن



النسج المسمى



أنتوا ازاي تغلوا النعناع كده .. كده غلط
النعناع بيتعمل infusion مش decoction

@kareem ayyad




يا بني بطل بقي الهبل اللي بتقوله دا
انت صدقت إنك دكتور بجد ولا إيه



➤ COMMON METHODS OF PLANT EXTRACTION

- **Percolation:**

In this method, the plant material is subjected to a slow flow of fresh solvent at intervals until sufficient active ingredient is extracted.



Commercial scale (about 1 ton capacity) percolator



E. Soxhlet extraction:

- Soxhlet extraction is only required where the desired compound has a limited solubility in a solvent, and the impurity is insoluble in that solvent.
- If the desired compound has a high solubility in a solvent then a simple filtration can be used to separate the compound from the insoluble substance.
- The advantage of this system is that instead of many portions of warm solvent being passed through the sample, just one batch of solvent is recycled.
- This method cannot be used for thermolabile compounds as prolonged heating may lead to degradation of compounds

يستخدم
كمية مذبذب
ولعادة
لدوري

وهي الطريقة ما يتكرر استخدامها
في التحليلات الحساسة للحرارة لأن ذلك
قد يؤدي إلى تدهور المركبات.


Q36. Which of the following statements about Soxhlet extraction is TRUE?

- A. It is suitable for thermolabile compounds because no heating is involved.
- B. It is used only when the desired compound has limited solubility in the solvent.
- C. It requires many fresh portions of warm solvent to pass through the sample.
- D. It is mainly used for extracting volatile oils by steam distillation.
- E. It is preferred when the desired compound has high solubility and can be separated by simple filtration.

B

Soxhlet extraction

Extraction: Withdrawal of soluble constituents from crude drug by treatment with solvent



Concentration of filtrate to dryness


- The filtered plant part (liquid portion) must be reduced to paste-like, cakey, and finally powdered form through standard techniques.
- The essence of concentrating filtrate to dryness is to:
- [?] Calculate the active ingredient present after the extraction by calculation of percentage yield.
- [?] To obtain dose of plant extract in milligram or gram that can be further constituted in a known volume of vehicle for experimental animals dosing.

The enriched extract from maceration or extractors, concentrated by;

- [?] Feding into a water bath, oven, hot plate, heating mantle, ...etc to produce a thick concentrated extract free from solvent.
- [?] Or the concentration could be done using rotary evaporator in order to remove excess solvents from samples by applying heat to a rotating vessel at a reduced pressure.
- [?] A freeze dryer or lyophilizer can also be used to concentrate and dry the filtrate extract (only aqueous) at reduced temperature

➤ COMMON METHODS OF CONCENTRATING FILTRATE TO DRYNESS

كمية ال solvent كبيرة يستخدم



Rotatory evaporator



Heating mantle



Freeze dryer (Lyophilizer)

GRAPHICAL OUTLINE



By Erhirre Earnest O.