

PHYSIOLOGY

MORPHINE ACADEMY

MORPHINE ACADEMY





PHYSIOLOGY

FACULTY OF PHARMACEUTICAL SCIENCES DR. AMJAAD ZUHIER ALROSAN

LECTURE 2, PART (1): CELLULAR LEVEL OF ORGANIZATION

Objectives

1. Discuss cellular level of organization. (Continue the previous lecture)---Part (1)

2. Describe transport processes of solutes and water .---- Part (2)


(Pages 60-84 of the reference)

THE ORGANELLES

Organelles are specialized structures within the cell that have characteristic shapes, and they perform specific functions in cellular growth, maintenance, and reproduction. They often cooperate to maintain homeostasis.

بتعانوا عشان يحافظوا على التوازن الداخلي

Organelles of the Cell




THE ORGANELLES

- 1. Centrosome.
- 2. Cilia and Flagella.
- 3. Ribosomes.
- 4. Endoplasmic Reticulum.
- 5. Golgi Complex.
- 6. Lysosomes.
- 7. Peroxisomes.
- 8. Proteasomes.
- 9. Mitochondria.
- 10. Nucleus.

Centrosome

- Is located near the nucleus and it consists of two components: a pair of centrioles and pericentriolar material.
- The centrioles are cylindrical structures (microtubules). Surrounding the centrioles is pericentriolar material which contains the tubulin complexes.
- These complexes organize centers for growth of the mitotic spindle (cell division).




Cilia

- Microtubules are the dominant components of cilia and flagella. Both are motile projections. However, cilia is present throughout the surface of a cell, but flagella is present at both the ends or all over the surface.
- Hair-like structure. بيكون شكلها زي الشعيرات
- The main function of motile cilia is to keep the airways clear of mucus and dust.

وتتحرك بشكل دائري rotational

Direction of motion




(b) Cilia

تنقسم الى نوعين

التي تتحرك motile cilia موجودة في الجهاز التنفسي respiratory system بتمسك المخاط الموجود على الرئتين عشان اطلعه للخارج

التي لا تتحرك non-motile cilia لما نؤخذ عن الحواس مسؤولة بشكل مباشر عن انتاج action potential

Flagella



(a) Flagella

Direction of motion

- Flagella are similar in structure to cilia but are typically much longer.
- Flagella usually move an entire cell (i.e. sperm).
- The motion of cilia is rotational, very fast moving. The motion of flagella is rotary movement in prokaryotes whereas it is bending movement in eukaryotes. Cilia beat in coordination واحدة تلو الأخرى بشكل or one after the other. Flagella beat independent of each other.

حركتها مستقلة عن بعضها البعض، أي كل سوط يتحرك بحرية دون تنسيق مع الآخر

Ribosomes

- Are the sites of protein synthesis.
- The name of these tiny structures reflects their high content of one type of ribonucleic acid (ribosomal RNA, or rRNA).
- "Scattered throughout cytoplasm": Some ribosomes are attached to the outer surface of the nuclear membrane and to an endoplasmic reticulum. Ribosomes are also located within mitochondria. Other ribosomes are "free" or unattached to other cytoplasmic structures.


يتم صناعة ribosomes من النواة nucluse من المادة الواثية DNA و هو مسؤول عن صناعة البر وتينات Amino Growing peptide acid chain Ribosome large subunit **tRNA** mRNA Ribosome small

subunit

Endoplasmic Reticulum

- The endoplasmic reticulum (ER) is a network of membranes in the form of flattened sacs or tubules.
- The ER extends from the nuclear envelope (membrane around the nucleus), to which it is connected and projects throughout the cytoplasm.
- Cells contain two distinct forms of ER (rough ER and smooth ER), which differ in structure and function.

Endoplasmic reticulum



Endoplasmic Reticulum

عليه

The outer surface of **rough ER** is studded with ribosomes (membrane-bound ribosomes). Proteins synthesized by ribosomes enter spaces within the ER for processing and sorting. Thus, rough ER produces secretory proteins, membrane proteins, and many organellar proteins.


وظیفتها انها لما یتم تصنیع البروتینات من ribosomse بتعمل علی تصنیف categorization وبترتبهم وکل مجموعة بروتینات بتعملهم تغلیف packaging بطلعهم بکیس vesicles وبروحوا علی golgi complex

• Unlike rough ER, smooth ER does not have ribosomes on its membrane. However, smooth ER contributes to lipids and carbohydrates synthesis, making phospholipids bilayer for cell membranes as well as it stores and releases calcium ions that trigger contraction in muscle cells.

وظيفتها تصنيع الكربو هيدرات والدهون واهم وظيفة هي كما أنه يخزن ويطلق أيونات الكالسيوم التي ً تحفز الانقباض في خلايا العضلات.

Golgi Complex

 Most of the proteins synthesized by ribosomes attached to rough ER are ultimately transported to other regions of the cell (i.e. Golgi complex).



Golgi Complex

Functions of Golgi Complex:

e two phases في عنا مرحلتين two phases في عنا مرحلتين cis phase trans phase

والمواد ممكن تفوت من cis phase وتطلع من trans phase والعكس كمان بس اهم اشي

- 1. Modifies, sorts, packages, and transports proteins received from the rough ER (through face or cis entry). Enzymes in the middle site of the Golgi complex modify the proteins to form glycoproteins, glycolipids, and lipoproteins.

 and lipoproteins.


 proteasomes وعلى الإغلب بتروح على مركب يسمى lipoprotine
- 2. Forms secretory vesicles that discharge processed proteins via exocytosis (through exist or trans face) into extracellular fluid and forms transport vesicles that carry molecules to other organelles, such as lysosomes. المعالّجة عن طريق endocytosis الى خارج الخلية، ويُشكّل حويصلات إفرازية تُفرِّغ البروتينات المعالّجة عن طريق endocytosis المحالة عن عضيات أخرى مثل lysosomes

Lysosomes

ممكن يحلل خليه كاملة او عضيات اوفيروسيات او يكمل تحليل مواد لم يتم اكتمال تحليها ويحتوي على انزيمات عديدة

- Lysosomes are membraneenclosed vesicles that form from the Golgi complex.
- They can contain as many as 60 kinds of powerful digestive and hydrolytic enzymes that can break down a wide variety of molecules.

LYSOSOME



Lysosomes

- Functions of lysosomes:
- 1. Digest substances (within the cell) that enter a cell via endocytosis and transport final products of digestion into cytosol.
- 2. Carry out autophagy, the digestion of worn-out organelles.
- 3. Implement autolysis, the digestion of an entire cell.
- 4. Accomplish extracellular digestion.

Peroxisomes

- Peroxisomes contain several oxidases, enzymes that can oxidize (remove hydrogen atoms from) various organic substances.
- They protect the cells from oxidative damage.
- They protect other parts of the cell from the toxic effects of potentially toxic compounds.
- Peroxisomes are very abundant in the liver, where detoxification of alcohol and other damaging substances occurs. توجد بشكل كبير في الكبد liver الذي يعتبر
- Peroxisomes also contain the enzyme catalase, which decomposes by-product of the oxidation reactions which is hydrogen peroxide (H₂O₂); thus, it decomposes hydrogen peroxide to water and oxygen. 2H₂O₂ Catalase

 $2H_{2}O + O_{2}$ يتم التخلص من H2O2 المادة جدًا سامة عن طريق انزيم catalase ويحلولها الى ماء و اكسجين




هذا الجزء الخارجي بمسك هذه البروتينات بعرف اذا هذه البروتينات الداخلي ويكسر ها ويحولها الى احماض امينية amino acid وممكن نستخدمها عشان انسوى بروتينات ثانية اذا بدنا

التخلص من البروتينات الموجودة في cytosol

- Cytosolic proteins require disposal at certain times in the life of a cell. Continuous destruction of unneeded, damaged, or faulty proteins is the function of tiny barrel-shaped structures consisting of four stacked rings of proteins around a central core called **proteasomes**. يحتوى على انزيمات تحلل البرو تينات فقط
- Proteasomes were so named because they contain proteases, enzymes that cut proteins into small peptides. Once the enzymes of a proteasome have chopped up a protein into smaller chunks, other enzymes then break down the peptides into amino acids, which can be recycled into new proteins.


26S proteasome



Mitochondria


- Mitochondria generate most of the ATP through aerobic (oxygen requiring) respiration, ("powerhouses" of the cell). The enzymes that catalyze the chemical reactions which are part of the aerobic phase of cellular respiration are located in the mitochondria.
- Active cells that use ATP at a high rate—such as those found in the muscles, liver, and kidneys—have a large number of mitochondria.
- A mitochondrion consists of an outer mitochondrial membrane and an inner mitochondrial membrane with a small fluid-filled space between them (mitochondrial matrix.).

Mitochondria Structural Features



Nucleus

- The nucleus is a spherical or oval-shaped structure that usually is the most prominent feature of a cell.
- Most cells have a single nucleus, although some, such as mature red blood cells, have none. Other types of cells have multiple nuclei.
- A double membrane called the <u>nuclear envelope</u> separates the nucleus from the cytoplasm. Many openings called <u>nuclear pores</u> extend through the nuclear envelope.
- The complex of DNA, proteins, and some RNA is called **chromatin**. The total genetic information carried in a cell, or an organism is its **genome**.



Nucleus

- Functions of nucleus:
- 1. Controls cellular structure.
- 2. Directs cellular activities.
- 3. Produces ribosomes in nucleoli.
- 4. Nuclear pores control the **movement of substances** between the nucleus and cytoplasm.

Membrane permeability:

- The permeability of the plasma membrane to different substances varies.
- The hydrophobic interior of the plasma membrane allows nonpolar molecules to rapidly pass through, but prevents passage of ions and large, uncharged polar molecules.
- Because water and urea are small polar molecules that have no overall charge, they can move from one gap (small gaps appear in the hydrophobic environment of the membrane's interior) to another until they have crossed the membrane without any assistance.
- Transmembrane proteins that act as channels and carriers (very selective) increase the plasma membrane's permeability to a variety of ions and uncharged polar molecules (need assistance).

Gradients across the Plasma Membrane:


- A <u>concentration gradient</u> is a difference in the concentration of a chemical from one place to another, such as from the inside to the outside of the plasma membrane. A substance will move "downhill," from where it is more concentrated to where it is less concentrated, to reach equilibrium.
- The plasma membrane also creates a difference in the distribution of positively and negatively charged ions between the two sides of the plasma membrane.

Gradients across the Plasma Membrane:

• The difference in electrical charges between two regions constitutes an <u>electrical</u> <u>gradient</u>. This termed the membrane potential. A positively charged substance will tend to move toward a negatively charged area, and a negatively charged substance will tend to move toward a positively charged area.

• The combined influence of the concentration gradient and the electrical gradient on movement of a particular ion is referred to as its *electrochemical gradient*.

الخلية لما تكون في حالة الراحة resting فرق الجهد voltage بين داخل الخلية وخارجها بيكون لازم يكون 70- يعني لما احكي electrical gradient هو gradient الي بحتاجه بحيث اوصل الخلية لوضع الراحة باستثناء except العضلة القلبية 20 وضع الراحة تبعها 90-



اما اذا المادة مشحونة وغير متعادلة فبدنا ننتبه concentration gradient وكمان electrical gradient

يعني المواد الي شحنتها موجبة رح تفضل تروح على المنطقة المشحونة بالشحنة السالبة والعكس صحيح

TRANSPORT ACROSS THE PLASMA MEMBRANE

- Substances generally move across cellular membranes via **transport processes** that can be classified as passive or active, depending on whether they require cellular energy.

ATP طاقة ATP

لانه تتنقل المواد من المكان التركيز العالى الى المكان التركيز قليل

- In passive processes, a substance moves down its concentration or electrical gradient to cross the membrane using only its own kinetic energy (energy of motion).

لانه تنتقل المواد من المكان التركيز القليل الى المكان التركيز العالي

- In active processes, cellular energy (in the form of adenosine triphosphate (ATP)) is used to drive the substance "uphill" against its concentration or electrical gradient.

TRANSPORT ACROSS THE PLASMA MEMBRANE

وعملية دخول المواد الى الخلية تسمى exocytosis وعملية خروج المواد من الخلية تسمى

Another way that some substances may enter and leave cells is an active process in which tiny, spherical membrane sacs referred to as vesicles are used. Examples include endocytosis, in which vesicles detach from the plasma membrane while bringing materials into a cell, and exocytosis, the merging of vesicles with the plasma membrane to release materials from the cell.

PASSIVE PROCESSES

Diffusion
Osmosis

الانتشار

- **Diffusion** is a passive process in which the random mixing of particles (solutes, the dissolved substances, and the solvent) in a solution occurs because of the particles' kinetic energy. الطاقة الحركية

- If a particular solute is present in high concentration in one area of a solution and in low concentration in another area, solute molecules will diffuse toward the area of lower concentration—they move down their concentration gradient (the solution after diffusion is said to be at equilibrium).

with a particular solute is present in high concentration in one area of a solute will another area, solute molecules will diffuse toward the area of lower concentration—they move down their concentration gradient (the solution after diffusion is said to be at equilibrium).

PASSIVE PROCESSES

- Substances may also diffuse through a membrane, if the membrane is permeable to them. Several factors influence the diffusion rate of substances across plasma membranes:

التناسب طردي direct

1. Steepness of the concentration gradient (the greater the difference in concentration between the two sides of the membrane, the higher is the rate of diffusion). يعني كل ما كان فرق التركيز والجهد بين داخل وخارج الخليه كل direct ما كان الانتقال والانتشار أسرع

2. <u>Temperature</u> (the higher the temperature, the faster the rate of diffusion).

التناسب عكسي Inverse

3. Mass of the diffusing substance (the larger the mass of the diffusing particle, the slower its diffusion rate).

التناسب طردي direct

4. <u>Surface area</u> (the larger the membrane surface area available for diffusion, the faster is the diffusion rate). هي المساحة المتاحة على الغشاء التي يمكن للجزيئات أن تمرّ من خلالها.

التناسب عكسي Inverse

5. <u>Diffusion distance</u> (The greater the distance over which diffusion must occur, the longer it takes).

هي المسافة اللي لازم الجزيئات تمشيها عشان تنتقل من جهة إلى أخرى (عبر الغشاء أو في السائل)

DIFFUSION


1- Simple diffusion

a substance moves across the lipid bilayer of the plasma membrane without the help of membrane transport proteins

2- Facilitated diffusion

channel-mediated facilitated diffusion, and carrier-mediated facilitated diffusion

A substance moves across the lipid bilayer aided by a channel protein or a carrier protein



DIFFUSION

Examples of simple diffusion

Nonpolar, hydrophobic molecules include oxygen, carbon dioxide, and nitrogen gases; fatty acids; steroids; and fat-soluble vitamins (A, D, E, and K) as well as small, uncharged polar molecules such as water, urea, and small alcohols.

Examples of facilitated diffusion

Too polar or highly charged solutes.

FACILITATED DIFFUSION

1. Channel- mediated facilitated diffusion

• Most membrane channels are <u>ion channels</u> integral transmembrane proteins that allow passage of <u>small</u>, <u>inorganic ions</u> that are too hydrophilic to penetrate the <u>nonpolar</u> interior of the lipid bilayer.

• Some gated channels randomly alternate between the open and closed positions; others are regulated by chemical or electrical changes inside and outside the cell. When the gates of a channel are open, ions diffuse into or out of cells, down their electrochemical gradients.

FACILITATED DIFFUSION

2. Carrier- mediated facilitated diffusion

- The solute binds more often to the carrier on the side of the membrane with a higher concentration of solute and is released on the other side after the carrier undergoes a change in shape.
- Once the concentration is the same on both sides of the membrane, solute molecules bind to the carrier on the cytosolic side and move out to the extracellular fluid.
- Substances that move across the plasma membrane by carrier mediated facilitated diffusion include glucose, fructose, galactose, and some vitamins.

OSMOSIS

انتقال الماء من تركيز الماء فيه عالي الى المكان تركيز الماء فيه قليل انتقال الماء من المكان تركيز sloute فيه قليل الى المكان تركيز فيه sloute فيه عالي

- Osmosis is a type of diffusion that occurs only when a membrane is permeable to water but is not permeable to certain solutes.

- In living systems, the solvent is water, which moves by osmosis across plasma membranes from an area of higher water concentration to an area of lower water concentration. In other words, water moves through a selectively permeable membrane from an area of lower solute concentration to an area of higher solute concentration.

OSMOSIS

التركيز داخل الخلية يساوي التركيز خارج الخلية

Any solution in which a cell maintains its normal shape and volume is an **isotonic solution**. The concentrations of solutes that cannot cross the plasma membrane are the same on both sides of the membrane in this solution.


تركيز منخفض خارج الخليه

- A different situation results if the cells are placed in a hypotonic solution, a solution that has lower concentration of solutes than the cytosol inside the cells. In this case, water molecules enter the cells faster than they leave, causing the cells to swell and eventually to burst (lysis).

OSMOSIS

تركيز مرتفع خارج الخلية

- A hypertonic solution has a higher concentration of solutes than does the cytosol inside cells. In such a solution, water molecules move out of the cells faster than they enter, causing the cells to shrink. Such shrinkage of cells is called crenation.



ACTIVE PROCESSES

- Two sources of cellular energy can be used to drive active transport:
- 1. Energy obtained from hydrolysis of adenosine triphosphate (ATP) is the source in **primary active transport**.
- 2. Energy stored in an ionic concentration gradient is the source in secondary active transport.


- Carrier proteins that mediate primary active transport are often called pumps.

ACTIVE PROCESSES

The most prevalent primary active transport mechanism expels sodium ions from cells and brings potassium ions in (sodium-potassium pump). All cells have thousands of sodium-potassium pumps in their plasma membranes.

SODIUM-POTASSIUM PUMP The splitting of ATP provides The sodium-potassium The sodium ions are re-Release of the phosphate pump binds three sodium energy to change the shape leased to the outside of the allows the channel to revert ions and a molecule of ATP. of the channel. The sodium membrane, and the new to its original form, releasions are driven through the shape of the channel ing the potassium ions on channel. allows two potassium ions the inside of the membrane. to bind.

ACTIVE PROCESSES



TRANSPORT IN VESICLES

- During **endocytosis**, materials move into a cell in a vesicle formed from the plasma membrane.
- In <u>exocytosis</u>, materials move out of a cell by the fusion with the plasma membrane of vesicles formed inside the cell.
- Both endocytosis and exocytosis require energy supplied by ATP. Thus, **transport in vesicles is an active process.**


- The three types of endocytosis:
- 1. Receptor-mediated endocytosis.
- 2. Phagocytosis.
- 3. Bulk-phase endocytosis.

RECEPTOR-MEDIATED ENDOCYTOSIS

زي ما حكينا من قبل انه بكون على المستقبل موقع نشط عشان يرتبط مع (المادة ligands زي LDL,والمضادات الحيوية antibiotics,وبروتين في الدم transferrin،الهرمونات)

- Cells take up cholesterol-containing low-density lipoproteins (LDLs), transferrin (an iron-transporting protein in the blood), some vitamins, antibodies, and certain hormones by receptor-mediated endocytosis.

يرتبط هذا المستقبل (reseptor LDL) مع مادة LDL وياخذوا جزء من LDL وياخذوا الخيه وبعملوا زي غطاء coat وبدخلوا داخل الخليه وبصير زي اندماج fusion مع lysosomes الي بيحطم المادة بس لما يخرجوا ، المستقبل plasma membran برجع الى reseptor برجع الى LDL جديد وبطلعوا اي اشي ما دها الخلية



PHAGOCYTOSIS


- **Phagocytosis** is a form of endocytosis in which the cell engulfs large solid particles, such as worn-out cells, whole bacteria, or viruses.

- Two main types of phagocytes are <u>macrophage</u>s, located in many body tissues, and <u>neutrophils</u>, a type of white blood cell.

PHAGOCYTOSIS

- Any undigested materials in the phagosome remain indefinitely in a vesicle called a residual body. The residual bodies are then either secreted by the cell via exocytosis or they remain stored in the cell.

الاقدام الكاذبة pseudopods داخلها يوجد reseptor ويرتبط فيه phagosome وبنفس الطريقة تدخل الخلية وبصير اسمها microbe وبصير تحليل digestion لهذا



BULK-PHASE ENDOCYTOSIS (PINOCYTOSIS)

- Bulk-phase endocytosis is a form of endocytosis in which tiny droplets of extracellular fluid are taken up (all solutes dissolved in the extracellular fluid are brought into the cell.).


- No receptor proteins are involved.

النوع الوحيد من endocytosis الي ما له reseptor

- Bulk-phase endocytosis occurs in most cells, especially absorptive cells in the intestines and kidneys.
- The resulting smaller molecules leave the lysosome to be used elsewhere in the cell.



water + solute



EXOCYTOSIS


- In contrast with endocytosis, which brings materials into a cell, exocytosis releases materials from a cell.

- All cells carry out exocytosis, but it is especially important in two types of cells: (1) secretory cells that liberate digestive enzymes, hormones, mucus, or other secretions and (2) nerve cells that release substances called neurotransmitters.

EXOCYTOSIS

- In some cases, wastes are also released by exocytosis. During exocytosis, membrane-enclosed vesicles called **secretory vesicles** form inside the cell, fuse with the plasma membrane, and release their contents into the extracellular

fluid.



انه في عمليه endocytosis بخسر جزء من plasma membran عشان لما تدخل المادة الى الخلية لازم تكون محاطة بغشاء ويتم تعويض هذه الخسارة عن طريق عملية exocytosis

Segments of the plasma membrane lost through endocytosis are recovered or recycled by exocytosis. The balance between endocytosis and exocytosis keeps the surface area of a cell's plasma membrane relatively constant.

TRANSCYTOSIS

- Vesicles undergo endocytosis on one side of a cell, move across the cell, and then undergo exocytosis on the opposite side.

- Transcytosis occurs most often across the endothelial cells that line blood vessels and is a means for materials to move between blood plasma and interstitial fluid.

ر المشيمة الاجسام المضادة

- For instance, when a woman is pregnant, some of her antibodies cross the placenta into the fetal circulation via transcytosis.

الدورة الدموية للجنين



THANK YOU

AMJADZ@HU.EDU.JO