


<u>ج</u>ان الدُّفعات

# PHYSIOLOGY

MORPHINE ACADEMY

MORPHINE ACADEMY



LECTURE 2, PART (1): CELLULAR LEVEL OF ORGANIZATION

# Objectives

1. Discuss cellular level of organization. (Continue the previous lecture)---Part (1)

2. Describe transport processes of solutes and water .---- Part (2)

(Pages 60-84 of the reference)

- **Cytoplasm** consists of all the cellular contents between the plasma membrane and the nucleus.

السيتوبلازم يتكون من جميع المحتويات الخلوية الواقعة بين النواة والعشاء البلازمي

- It has two components:
- (1) the cytosol, is the fluid portion of the cytoplasm that surrounds organelles.
- (2) organelles, are tiny structures that perform different functions in the cell.

العضيات:هي تراكيب صغيرة تقوم بوظائف متعددة في الخلية

## - Cytosol:

السيتوسول يحتوي كمية كبيرة من الماء بنسبة 75-90% ويحتوي العديد من المواد المذابة والمعرفة مثل :سكر الغلوكوز و الأحماض الامينية والدهنية والبروتين والليبيدات و ATP والفضلات


1. Contains 75–90% water plus various dissolved and suspended components (i.e. glucose, amino acids, fatty acids, proteins, lipids, ATP, and waste products).

2. It is the site of many chemical reactions required for a cell's existence (i.e. glycolysis, maintenance of cell structures and for cell growth).

السيتوسول هو موقع للكثير من التفاعلات الكيميائية من هدم وتكسير مكونات وصناعة المواد التي تحتاجها الخلية ك:تحليل سكر الغلوكوز والحفاظ على وجود الخلية ونموها وتركيبها

- The cytoskeleton is a network of protein filaments that extends throughout the cytosol (microfilaments, intermediate filaments, and microtubules).

and microtubules).



### - Microfilaments: 3 to 6 nm diameter

1. Are the thinnest elements of the cytoskeleton.

الفع عنفسر في، لم يكل الخلوب المحلف المحلوب المحلوب المحلوب الميوسين

- 2. They are composed of the proteins actin and myosin.
- 3. They have two general functions: help generate movement (muscle contraction, cell division, and cell locomotion) and provide mechanical support (basic strength and shapes of cells).

### - Intermediate filaments:

- 1. Are thicker than microfilaments but thinner than microtubules.
- 2. They are found in parts of cells subject to mechanical stress.
- 3. They help stabilize the position of organelles such as the nucleus and help attach cells to one another.

### - Microtubules:


- 1. Are the largest of the cytoskeletal components.
- 2. They are composed mainly of the protein tubulin.
- 3. They help determine cell shape.
- 4. They also function in the movement of organelles.

اهم الخصائص لل :actin filament هو أنه أكبر مكونات الهيكل الخلوي ويتكون بشكل رئيس من بروتين ال tubulin وتعمل على (تحديد شكل الخلية ) وحركة العضيات في السيتوبلازم

#### THE ORGANELLES

Organelles are specialized structures within the cell that have characteristic shapes, and they perform specific functions in cellular growth, maintenance, and reproduction. They often to maintain cooperate homeostasis الموازن egrili brium

# Organelles of the Cell



العضيات هي تراكيب مميزة في الخلية وهي أحد مكونات السيتوبلازم /تقوم بوظائف محددة كالنمو الخلوي والإصلاح والتكاثر وهي تشارك في الحفاظ على توازن الخلية


## THE ORGANELLES


- 1. Centrosome.
- 2. Cilia and Flagella.
- المطلود: حفا لقه كالوظيفة كالوقع

- 3. Ribosomes.
- 4. Endoplasmic Reticulum.
- 5. Golgi Complex.
- 6. Lysosomes.
- 7. Peroxisomes.
- 8. Proteasomes.
- 9. Mitochondria.
- 10. Nucleus.

## Centrosome

- Is located near the nucleus and it consists of two components: a pair of centrioles and pericentriolar material.
- The centrioles are cylindrical structures (microtubules). Surrounding the centrioles is pericentriolar material which contains the tubulin complexes.
- These complexes organize centers for growth of the mitotic spindle (cell division).






## Cilia


- Microtubules are the dominant components of cilia and flagella. Both are motile projections. However, cilia is present throughout the surface of a cell, but flagella is present at both the ends or all over the surface.
- Hair-like structure.
- The main function of motile cilia is to keep the airways clear of mucus and dust.

cilia: هي شعيرات متحركة توجد بشكل رئيس في الجهاز التنفسي لإزالة المخاط والغبار وإخراجه خارج الجسم



(b) Cilia

# Flagella



حركة ال cilia متناسقة أو وحدة ورا وحدة اي باتجاه واحد/أما ال flagella فهي حركتها مستقلة تكون في حقيقيات النوى حركة كموجة waving وفي بدائيات النوى حركته دورانية (a) Flagella

#### **Direction of motion**




- Flagella are similar in structure to cilia but are typically much longer.
- Flagella usually move an entire cell (i.e. sperm).
  - The motion of cilia is rotational, very fast moving. The motion of flagella is rotary movement in prokaryotes whereas it is bending movement in eukaryotes. Cilia beat in coordination or one after the other. Flagella beat independent of each other.

## Ribosomes

- Are the sites of protein synthesis.
- The name of these tiny structures reflects their high content of one type of ribonucleic acid (ribosomal RNA, or rRNA).

Location of vibosome:-

"Scattered throughout cytoplasm": Some ribosomes are attached to the outer surface of the nuclear membrane and to an endoplasmic reticulum. Ribosomes are also located within mitochondria. Other ribosomes are "free" or unattached to other cytoplasmic structures.



المختصر:الريبوسوم هو مصنع البروتين وسبب تسميته بأنه يحتوي على نسبة كبيرة من الحمض النووي الرايبوسومي rRNA / يوجد الريبوسوم على سطح الغشاء النواة /وعلى الشبكة الاندوبلازمية/ الميتوكندريا/وحرة غير مرتبطة في السيتوبلازم

# Endoplasmic Reticulum

 The endoplasmic reticulum (ER) is a network of membranes in the form of flattened sacs or tubules.


الشبكة الاندوبلازمية هي شبكة من الأغشية على شكل اكياس وأنابيب مستوية

• The ER extends from the nuclear envelope (membrane around the nucleus), to which it is connected and projects throughout the cytoplasm.

تمتد الشبكة الاندوبلازمية من الغشاء النووي المتصلة به وتنطلق خلال السيتوبلازم

 Cells contain two distinct forms of ER (rough ER and smooth ER), which differ in structure and function.

#### Endoplasmic reticulum



الخلايا تحتوي على نوعين من الشبكة الاندوبلازمية وهما الخشنة والنعمة وكلاهما يختلف بالترحيب والوظيفة

# Endoplasmic Reticulum

## (هم شي نفرن المواد ١

The outer surface of **rough ER** is studded with ribosomes (membrane-bound ribosomes). Proteins synthesized by ribosomes enter spaces within the ER for processing and sorting. Thus, rough ER produces secretory proteins, membrane proteins, and many organellar proteins.

Illumpater of the product of the product of the product of the product of the proteins of the product of the pro


الشبكة الخشنة تكون الرايبوسومات منغمسة في غشائها وتعمل على إنتاج البروتين الذي يدخل المساعدة داخل الشبكة الشبكة الاندوبلازمية ويحدث للبروتين معالجة وفرز /الشبكة تعمل على معالجة البروتين ليكوّن عضيات بروتيني كالبروتينات الافرازية والغشائية،...

• Unlike rough ER, smooth ER does not have ribosomes on its membrane. However, smooth ER contributes to lipids and carbohydrates synthesis, making phospholipids bilayer for cell membranes as well as it stores and releases calcium ions that trigger contraction in muscle cells.

# Golgi Complex

 Most of the proteins synthesized by ribosomes attached to rough ER are ultimately transported to other regions of the cell (i.e. Golgi complex).

البروتين الذي أنتج في الشبكة الاندوبلازمية ينتقل لجهاز جولجي في النهاية



# Golgi Complex


- لمختصر يعمل جهاز جولجي على التعديل وفرز وتغليف البروتين وإضافة تعديلات عليه لإنتاج مواد مثل /glycoprotein Complex والمحتصر يعمل جهاز جولجي على التعديل وفرز وتغليف البروتين وإضافة تعديلات عليه لإنتاج مواد مثل /glycoprotein Functions of Golgi Complex المحتال المعالج إلى السائل الخلوي ومن ثم تنشأ حويصلة ناقلة لتنقل البروتين المعالج إلى glycoprotein المحتال المحتال على السائل الخلوي ومن ثم تنشأ حويصلة ناقلة لتنقل البروتين المعالج إلى المحتال الم
- 1. Modifies, sorts, packages, and transports proteins received from the rough ER (through face or cis entry). Enzymes in the middle site of the Golgi complex modify the proteins to form glycoproteins, glycolipids, and lipoproteins.

2. Forms secretory vesicles that discharge processed proteins via exocytosis (through exist or trans face) into extracellular fluid and forms transport vesicles that carry molecules to other organelles, such as lysosomes.

# Lysosomes

- Lysosomes are membraneenclosed vesicles that form from the Golgi complex.
- They can contain as many as 60 kinds of powerful digestive and hydrolytic enzymes that can break down a wide variety of molecules.

LYSOSOME



يمكن أن تحتوي على ما يصل إلى 60 نوعًا من الإنزيمات الهاضمة والتحليلية القوية، القادرة على تفكيك مجموعة واسعة من الجزيئات.

# Lysosomes

- Functions of lysosomes:
- 1. Digest substances (within the cell) that enter a cell via endocytosis and transport final products of digestion into cytosol.

عمل هيم عبر الانزيمات للمواد التي يدفل الخلية عبر bostosbus عمر تنقل نواتج الهام الحي الدون عاده b المناق التي

- 2. Carry out autophagy, the digestion of worn-out organelles.
- 3. Implement autolysis, the digestion of an entire cell. bacteria
- 4. Accomplish extracellular digestion.

ليكا ي كن بنعوا عيلمد كالم إ

## Peroxisomes

- Peroxisomes contain several oxidases, enzymes that can oxidize (remove hydrogen atoms from) various organic substances.
- ير له على النيموسوم يحتوي على الزيمات الأكسدة التي تعمل على إزالة الهيدروجين من المواد فبذلك تعمل على حماية الخلية من الضرر

  They protect the cells from oxidative damage.

  They protect the cells from oxidative damage.
- They protect other parts of the cell from the toxic effects of potentially toxic compounds.
- Peroxisomes are very abundant in the liver, where detoxification of alcohol and other damaging substances occurs.
- Peroxisomes also contain the enzyme catalase, which decomposes by-product of the oxidation reactions which is hydrogen peroxide ( $H_2O_2$ ); thus, it decomposes hydrogen peroxide to water and oxygen.

ليعة ي على الرامي ( معدما معله ع) الذي يعمل الله كم الله نوب من تناعلات الكاتحسر فيحل و Ho الى ماء و اكسجين

## Proteasomes


لترصير البير, تينات المثالثة

• Cytosolic proteins require disposal at certain times in the life of a cell. **Continuous destruction of unneeded, damaged, or faulty proteins** is the function of tiny barrel-shaped structures consisting of four stacked rings of proteins around a central core called **proteasomes**.

يحتوي على الزهائ لتعطيم البردين المعلوانة مكونة من (١٧) حلق عمواهة

Proteasomes were so named because they contain proteases, enzymes that cut proteins into small peptides. Once the enzymes of a proteasome have chopped up a protein into smaller chunks, other enzymes then break down the peptides into amino acids, which can be recycled into new proteins.


## 26S proteasome



## Mitochondria


- Mitochondria generate most of the ATP through aerobic (oxygen requiring) respiration, ("powerhouses") of the cell). The enzymes that catalyze the chemical reactions which are part of the aerobic phase of cellular respiration are located in the mitochondria. هر العقب الزي من العالمة عبر المناس الموالي و لد به الاترس الموالي و لد به الاترس الموالي و لد به الاترس الموالي و لد به المناسة عبر المناسة عبر المناسة عبر المناسة عبر المناسة عبر المناسة المناسة عبر المناسة المناسة عبر المناسة المناسة عبر المناسة عبر المناسة المناسة عبر المناسة المناسة عبر المناسة المن
- Active cells that use ATP at a high rate—such as those found in the muscles, liver, and kidneys—have a large number of mitochondria.
- A mitochondrion consists of an outer mitochondrial membrane and an inner mitochondrial membrane with a small fluid-filled space between them (mitochondrial matrix.).

#### Mitochondria Structural Features



## Nucleus

- The nucleus is a spherical or oval-shaped structure that usually is the most prominent feature of a cell.
- Most cells have a single nucleus, although some, such as mature red blood cells, have none. Other types of cells have multiple nuclei.
- A double membrane called the <u>nuclear envelope</u> separates the nucleus from the cytoplasm. Many openings called <u>nuclear pores</u> extend through the nuclear envelope.
- The complex of DNA, proteins, and some RNA is called **chromatin**. The total genetic information carried in a cell, or an organism is its **genome**.



# Nucleus


## Functions of nucleus:

- 1. Controls cellular structure.
- 2. Directs cellular activities.
- 3. Produces ribosomes in nucleoli.

آلتمكم مي تولي الحديد
 أب من طعل انشطة الخديد
 أب من طعل انشطة الخديد
 أب من طعل انشطة الخديد
 أب من الرايبو لموم في المؤلفيد
 أب لديها فجوان في غشائها تتكم مني مرو , المواد

بن النواه والسية بلازع.

4. Nuclear pores control the **movement of substances** between the nucleus and cytoplasm.




DR. AMJAAD ZUHIER ALROSAN

LECTURE 2, PART (2): TRANSPORT PROCESSES OF SOLUTES AND WATER

## Membrane permeability:

#### نفاذ وست الفشاء للعواد

• The permeability of the plasma membrane to different substances varies.



- The hydrophobic interior of the plasma membrane allows nonpolar molecules to rapidly pass through, but prevents passage of ions and large, uncharged polar molecules.

  الجزيات العيرمشجانة والكبيرة
- Because water and urea are small polar molecules that have no overall charge, they can move from one gap (small gaps appear in the hydrophobic environment of the membrane's interior) to another until they have crossed the membrane without any assistance.

معنوع تقر حذ المفتاد بامتثناء ال معلم مهم معنوع تقر حذ المفتاد بامتثناء ال معلم مهم مهم المعنودة في و سط غشاء الذليم.

• Transmembrane proteins that act as channels and carriers (very selective) increase the plasma membrane's permeability to a variety of ions and uncharged polar molecules (need assistance).

(need assistance).

#### **Gradients across the Plasma Membrane:**

• A <u>concentration gradient</u> is a difference in the concentration of a chemical from one place to another, such as from the inside to the outside of the plasma membrane. A substance will move "downhill," from where it is more concentrated to where it is less concentrated, to reach equilibrium.

تدرج التركيز :هو اختلاف تراكيب المواد الكيميائية كالايونات على جانبي الغشاء

• The plasma membrane also creates a difference in the distribution of positively and negatively charged ions between the two sides of the plasma membrane.

downhill: with concentration gradient... from Large concentration to less concentration

احي من المنطقة الماعل تركيز الى الاقل توكيز الي ح قارع التوكيز


کے الفشاء بیعل منے بن الرّا لیز علی جانبیت ر معکن یجی دؤال ۲۰۰۰ نام عام

#### **Gradients across the Plasma Membrane:**

• The difference in electrical charges between two regions constitutes an <u>electrical gradient</u>. This termed the membrane potential. A positively charged substance will tend to move toward a negatively charged area, and a negatively charged substance will tend to move toward a positively charged area.

و بيعل عبد للخديث ، وبد ذلا الايون د لميل العركة ذعو الايون • و العكس كذلا ،

• The combined influence of the concentration gradient and the electrical gradient on movement of a particular ion is referred to as its electrochemical gradient.



# TRANSPORT ACROSS THE PLASMA MEMBRANE

المواد تعبر الصناء الخذي بعملية الانتقال

- Substances generally move across cellular membranes via transport processes that can be classified as passive or active, depending on whether they require cellular energy.

  \*\*Transport\*\* process\*\*

  \*\*Transport\*\*

  \*\*Tran
- In passive processes, a substance moves down its concentration or electrical gradient to cross the membrane using only its own kinetic energy (energy of motion).
- In active processes, cellular energy (in the form of adenosine triphosphate (ATP)) is used to drive the substance "uphill" against its concentration or electrical gradient.

# TRANSPORT ACROSS THE PLASMA MEMBRANE

active Evansport ا هم معلومة الله عنوontosis و molomtosis من أنواع ال

نستخدم المويصلات (۱۰۵۱ (۱۰۵۱ الاواد تعريف المريصة

Another way that some substances may enter and leave cells is an active process in which tiny, spherical membrane sacs referred to as vesicles are used. Examples include endocytosis, in which vesicles detach from the plasma membrane while bringing materials into a cell, and exocytosis, the merging of vesicles with the plasma membrane to release materials from the cell.

# PASSIVE PROCESSES

without extlutor every

facilitated diffusion 122

الانتشار

Diffusion

- **Diffusion** is a passive process in which the random mixing of particles (solutes, the dissolved substances, and the solvent) in a solution occurs because of the particles' kinetic energy.

والمزيب لهاطامة حرحية خاصة تحفلها لمرف الفشاء برون بعوريه

- If a particular solute is present in high concentration in one area of a solution and in low concentration in another area, solute molecules will diffuse toward the area of lower concentration—they move down their concentration gradient (the solution after diffusion is said to be at equilibrium).

## PASSIVE PROCESSES

جمیع الوامل تؤثر فردیا ما عدا مناعاته لی ما عدا مناعاته لی

- Substances may also diffuse through a membrane, if the membrane is permeable to them. Several factors influence the diffusion rate of substances across plasma membranes:

الاتحار

عوامل مؤثرة على صعدل عملية الانتشار

- 1. Steepness of the concentration gradient (the greater the difference in concentration between the two sides of the membrane, the higher is the rate of diffusion).
- 2. Temperature (the higher the temperature, the faster the rate of diffusion).
- 3. Mass of the diffusing substance (the larger the mass of the diffusing particle, the slower its diffusion rate).
- 4. <u>Surface area</u> (the larger the membrane surface area available for diffusion, the faster is the diffusion rate).
- 5. <u>Diffusion distance</u>(The greater the distance over which diffusion must occur, the longer it takes).

## **DIFFUSION**

## 1- Simple diffusion


a substance moves across the lipid bilayer of the plasma membrane without the help of membrane transport proteins

العبر دود ما عدة الزائل كما هي المو أد غير القطبية.

## 2- Facilitated diffusion

channel-mediated facilitated diffusion, and carrier-mediated facilitated facilitated

A substance moves across the lipid bilayer aided by a channel protein or a carrier protein



## DIFFUSION

Examples of simple diffusion

المواد التي تتبر احرى هي الرسيق

Nonpolar, hydrophobic molecules include oxygen, carbon dioxide, and nitrogen gases; fatty acids; steroids; and fat-soluble vitamins (A, D, E, and K) as well as small, uncharged polar molecules such as water, urea, and small alcohols.

## Examples of facilitated diffusion

Too polar or highly charged solutes.

عواد دان تعليقالية

### FACILITATED DIFFUSION

## 1. Channel- mediated facilitated diffusion

مخترتة الغثاء بالكان

- Most membrane channels are ion channels <u>integral</u> transmembrane proteins that allow passage of <u>small</u>, <u>inorganic</u> ions that are too <u>hydrophilic</u> to <u>penetrate</u> the nonpolar interior of the lipid bilayer.
- Some gated channels randomly alternate between the open and closed positions; others are regulated by chemical or electrical changes inside and outside the cell. When the gates of a channel are open, ions diffuse into or out of cells, down their electrochemical gradients.


## FACILITATED DIFFUSION

# 2. Carrier- mediated facilitated diffusion

The solute binds more often to the carrier on the side of the membrane with a higher concentration of solute and is released on the other side after the carrier undergoes a change in shape

- Once the concentration is the same on both sides of the membrane, solute molecules bind to the carrier on the cytosolic side and move out to the extracellular fluid.
- Substances that move across the plasma membrane by carrier mediated facilitated diffusion include glucose, fructose, galactose, and some vitamins.

السكويات الاحادية و بعف الفتي منيات



### **OSMOSIS**

Osmosis is a type of diffusion that occurs only when a membrane is permeable to water but is not permeable to certain solutes.

- In living systems, the solvent is water, which moves by osmosis across plasma membranes from an area of higher water concentration to an area of lower water concentration. In other words, water moves through a selectively permeable membrane from an area of lower solute concentration to an area of higher solute concentration.

تقريفين لعملية ال عاده osmosis

#### **OSMOSIS**


- Any solution in which a cell maintains its normal shape and volume is an **isotonic solution**. The concentrations of solutes that cannot cross the plasma membrane are the same on both sides of the membrane in this solution.

توكيز هتسلوي على جابني المخليك

- A different situation results if the cells are placed in a hypotonic solution, a solution that has lower concentration of solutes than the cytosol inside the cells. In this case, water molecules enter the cells faster than they leave, causing the cells to swell and eventually to burst (lysis).

#### **OSMOSIS**

- A hypertonic solution has a higher concentration of solutes than does the cytosol inside cells. In such a solution, water molecules move out of the cells faster than they enter, causing the cells to shrink. Such shrinkage of cells is called crenation.



### **ACTIVE PROCESSES**

عن طریت (حیل ۱۹۴۸ عن طریق نتر ای تولیز ۱ کایونی (عن طریق اختمال ایون مع ترریخ پینمتیل المکر مکس ترکیز

- Two sources of cellular energy can be used to drive active transport:
- 1. Energy obtained from hydrolysis of adenosine triphosphate (ATP) is the source in primary active transport.
- 2. Energy stored in an ionic concentration gradient is the source in secondary active transport.

- Carrier proteins that mediate primary active transport are often called

pumps


الستخرية نهزا المستخرية نهزا النوع الممهاطنخة


#### **ACTIVE PROCESSES**


الاقة المسادة لنتل مهم المجمع

The most prevalent primary active transport mechanism expels sodium ions from cells and brings potassium ions in (sodium-potassium pump). All cells have thousands of sodium-potassium pumps in their plasma membranes.


#### SODIUM-POTASSIUM PUMP








The splitting of ATP provides energy to change the shape of the channel. The sodium ions are driven through the channel.




The sodium ions are released to the <u>outside of the</u> membrane, and the new shape of the channel allows two potassium ions to bind.



Release of the phosphate allows the channel to revert to its original form, releasing the potassium ions on the inside of the membrane.

# **ACTIVE PROCESSES**



## TRANSPORT IN VESICLES

النعل اللي يتم من خلال الحولهلات يسمى
Transcytosis

- During <u>endocytosis</u>, materials move into a cell in a v<u>esicle formed from the plasma</u> membrane.
- In <u>exocytosis</u>, materials move out of a cell by th<u>e fusio</u>n with the plasma membrane of vesicles formed inside the cell.
- Both endocytosis and exocytosis require energy supplied by ATP. Thus, transport in vesicles is an active process.

- The three types of endocytosis:
- 1. Receptor-mediated endocytosis.
- 2. Phagocytosis.
- 3. Bulk-phase endocytosis.——>

ىلھمە


receptor

لأيحتاج هستقبق

# RECEPTOR-MEDIATED ENDOCYTOSIS

Cells take up cholesterolcontaining low-density
lipoproteins (LDLs)
transferrin (an irontransporting protein in
the blood), some
vitamins, antibodies, and
certain hormones by
receptor-mediated
endocytosis.

receptor mediated responsions endoustosis



#### **PHAGOCYTOSIS**

**Phagocytosis** is a form of endocytosis in which the cell engulfs large solid particles, such as worn-out cells, whole bacteria, or viruses.

فيروساء / بكيتربا / خلايا تالفة


Two main types of phagocytes are <u>macrophages</u>, located in many body tissues, and <u>neutrophils</u>, a type of white blood cell.

#### **PHAGOCYTOSIS**

Any undigested materials in the phagosome remain indefinitely in a vesicle called a residual body. The residual bodies are then either secreted by the cell via exocytosis or they remain stored in the cell.

محي اجزاء عير مهانومة سمى والمعمد المساه residual

و 10 ي الاحبام نتبي في الجم البلعمي او بمين للخلية طرد 10



# BULK-PHASE ENDOCYTOSIS (PINOCYTQSIS)

للعمة لكن لسائل نيد مواد دَحدا ع الخدة


Bulk-phase endocytosis is a form of endocytosis in which tiny droplets of extracellular fluid are taken up (all solutes dissolved in the extracellular fluid are brought into the cell.).

No receptor proteins are involved.

- Bulk-phase endocytosis occurs in most cells, especially absorptive cells in the intestines and kidneys. اکثر و بو دھا ہی ملکیوالعماء
- (The resulting smaller molecules) leave the lysosome to be used elsewhere in the cell.

لَفَادر المواد التي نتم هعنها في معنها في Lysosomes لتستضم في الخليد

# Bulk-phase Endocytosis



#### **EXOCYTOSIS**

- In contrast with endocytosis, which brings materials into a cell, exocytosis releases materials from a cell.


- All cells carry out exocytosis, but it is especially important in two types of cells: (1) secretory cells that liberate digestive enzymes, hormones, mucus, or other secretions and (2) nerve cells that release substances called neurotransmitters.



ا هم نوسين خلايا نقعل عفلية vxoontosis ا

#### **EXOCYTOSIS**

- In some cases, wastes are also released by exocytosis. During exocytosis, membrane-enclosed vesicles called **secretory vesicles** form inside the cell, fuse with the plasma membrane, and release their contents into the extracellular fluid.



Segments of the plasma membrane lost through endocytosis are recovered or recycled by exocytosis. The balance between endocytosis and exocytosis keeps the surface area of a cell's plasma membrane relatively constant.

لزلاد لازم کل ۱۱۵۶ مه همه دیسر مهه ۱۱۵۶ و ۱۵۵۸ مه کل ۱۱۵۴ مهمه دیستر مهد کمون میست الانادا کنون مب عملیت الانادا و در الانادا کنون مب عملیت الانادا و در الانادا کنون مب عملیت الانادا و در الانادا و در الانادا کنون مب عملیت الانادا و در ال

#### **TRANSCYTOSIS**

Vesicles undergo endocytosis on one side of a cell, move across the cell, and then undergo exocytosis on the opposite side.

عمليات الإدخال والاخراع الحنرك

المائل خارج الخلمة

تحدث منبة كبيرة في المحافظ في طبعة المه العلال له

- Transcytosis occurs most often across the endothelial cells that line blood vessels and is a means for materials to move between blood plasma and interstitial fluid.
- مرأة حاص - For instance, when a woman is pregnant, some of her antibodies cross the placenta into the fetal circulation via transcytosis. تَصَوعير المسينة الى الحين عبر دورته الموية



# THANK YOU

AMJADZ@HU.EDU.JO